This is a digital copy of a book that was preserved for generations on library shelves before it was carefully scanned by Google as part of a project to make the world's books discoverable online.

It has survived long enough for the copyright to expire and the book to enter the public domain. A public domain book is one that was never subject to copyright or whose legal copyright term has expired. Whether a book is in the public domain may vary country to country. Public domain books are our gateways to the past, representing a wealth of history, culture and knowledge that's often difficult to discover.

Marks, notations and other marginalia present in the original volume will appear in this file - a reminder of this book's long journey from the publisher to a library and finally to you.

Usage guidelines

Google is proud to partner with libraries to digitize public domain materials and make them widely accessible. Public domain books belong to the public and we are merely their custodians. Nevertheless, this work is expensive, so in order to keep providing this resource, we have taken steps to prevent abuse by commercial parties, including placing technical restrictions on automated querying.

We also ask that you:

+ *Make non-commercial use of the files* We designed Google Book Search for use by individuals, and we request that you use these files for personal, non-commercial purposes.

+ *Refrain from automated querying* Do not send automated queries of any sort to Google's system: If you are conducting research on machine translation, optical character recognition or other areas where access to a large amount of text is helpful, please contact us. We encourage the use of public domain materials for these purposes and may be able to help.

+ *Maintain attribution* The Google "watermark" you see on each file is essential for informing people about this project and helping them find additional materials through Google Book Search. Please do not remove it.

+ *Keep it legal* Whatever your use, remember that you are responsible for ensuring that what you are doing is legal. Do not assume that just because we believe a book is in the public domain for users in the United States, that the work is also in the public domain for users in other countries. Whether a book is still in copyright varies from country to country, and we can't offer guidance on whether any specific use of any specific book is allowed. Please do not assume that a book's appearance in Google Book Search means it can be used in any manner anywhere in the world. Copyright infringement liability can be quite severe.

About Google Book Search

Google's mission is to organize the world's information and to make it universally accessible and useful. Google Book Search helps readers discover the world's books while helping authors and publishers reach new audiences. You can search through the full text of this book on the web at `http://books.google.com/`

The University of Chicago

FOUNDED BY JOHN D. ROCKEFELLER

THE STRUCTURE AND DEVELOPMENT OF THE NEPHRIDIA OF ARENICOLA CRISTATA STIMPSON

A DISSERTATION

SUBMITTED TO THE FACULTY OF THE OGDEN GRADUATE SCHOOL OF SCIENCE IN CANDIDACY FOR THE DEGREE OF DOCTOR OF PHILOSOPHY

(DEPARTMENT OF ZOÖLOGY)

BY

RALPH S. LILLIE.

BUHR A
a39015 01801101 8b

CHICAGO

1905


1837
ARTES
VERITAS
SCIENTIA
LIBRARY OF THE
UNIVERSITY OF MICHIGAN
E PLURIBUS UNUM
SI QUAERIS PENINSULAM AMOENAM
CIRCUMSPICE

SCIENCE LIBRARY
QL
391
.P9
L73

The University of Chicago
FOUNDED BY JOHN D. ROCKEFELLER

THE STRUCTURE AND DEVELOPMENT OF THE NEPHRIDIA OF ARENICOLA CRISTATA STIMPSON

A DISSERTATION

SUBMITTED TO THE FACULTY OF THE OGDEN GRADUATE SCHOOL OF SCIENCE IN CANDIDACY FOR THE DEGREE OF DOCTOR OF PHILOSOPHY

(DEPARTMENT OF ZOÖLOGY)

BY
RALPH S. LILLIE

CHICAGO
1905

(Mittheilungen a. d. zoologischen Station zu Neapel. 17. Bd. 3. Heft. 1905.)

The Structure and Development of the Nephridia of Arenicola cristata Stimpson.

By

Ralph S. Lillie.

With Plates 22—25 and one Textfigure.

1. Introductory.

The nephridia of *Arenicola* were first suggested to me as a subject of anatomical and embryological investigation by Dr. C. O. WHITMAN, during the season of 1896 at the Marine Biological Laboratory of Wood's Hole. The work then begun was subsequently interrupted for a time; and the greater part of the work described in the following paper was completed during the years 1898—1901, at the Marine Biological Laboratory and in the Hull Zoölogical Laboratory of the University of Chicago.

I take pleasure in expressing my best thanks to Dr. WHITMAN for his continued interest and criticism. My thanks are also due to Dr. C. M. CHILD of the University of Chicago, for much helpful information, especially with reference to the early development; and to my brother, Professor F. R. LILLIE of the same University for constant and valuable assistance.

The nephridia of several species of the genus *Arenicola* have been at various times figured and briefly described in systematic or anatomical treatises; and in several instances have been made the subject of special investigation. CLAPARÈDE (1868) and FAUVEL (1899) have given brief descriptions of the nephridia of *A. Grubii* and *A. ecaudata* respectively. The greater number of such studies have been made on *A. marina*; investigations on the nephridia of this form have been made by COSMOVICI (1880), CUNNINGHAM (1887), BENHAM (1891, 1893), KYLE (1896); and more recently by GAMBLE &

ASHWORTH (1898, 1900), who in their two important papers on the anatomy and affinities of the Arenicolidae have added materially to our knowledge of these organs. The second paper of these authors (1900) contains a comparative account of the anatomy, relations and vascular supply of the nephridia of five species of the genus, viz: — *A. marina* Linn., *A. cristata* Stimpson, *A. Claparedii* Levinsen, *A. Grubii* Claparède and *A. ecaudata* Johnston; the authors have also given a summary of our present knowledge of the anatomy and histology of the adult nephridia, and have added a brief account of the condition of the organs in earlier developmental stages, so far as this has as yet been ascertained.

Our knowledge of the larval development of *Arenicola* has, however, until recently been too limited to admit of a detailed study of the mode of origin of the nephridia, and of the condition of these organs at different periods in the life-history. BENHAM (1893) and GAMBLE & ASHWORTH (1898, 1900) have contributed a few observations on the nephridia of certain post-larval stages. Their observations were made on relatively few specimens which in all cases were well-advanced in development. Accordingly, we have had as yet no detailed account of the origin, growth, and histological differentiation of the nephridia in *Arenicola*. One reason for this deficiency seems hitherto to have been the difficulty of obtaining the material for such an investigation. In my own studies, however, instead of depending upon the slow and precarious methods of digging and tow-net collecting, I have found it more practicable to rear the larvae artificially; in this manner it has been found possible to secure all the material necessary for a detailed examination of the larval development from the early free-swimming period up to a stage at which the adult characteristics are essentially complete.

The aim of the present study has been to complete so far as possible our knowledge of the origin and histological differentiation of the nephridia, and to determine to what extent the peculiarities of the adult nephridium gain an explanation through the conditions of larval development. The relations of the nephridia to the body-segmentation and to the septa, and the manner of development of the different portions of the organ — nephrostome, glandular portion and terminal vesicle (whether separately, or by differentiation from a single embryonic rudiment) — are matters that have received especial attention. There has also been added a brief account of the anatomy and histology of the adult nephridium, draw-

ing especial attention to a few peculiarities that have hitherto apparently escaped observation.

2. Technique.

The adult nephridia are best prepared for histological study by fixing in HERMANN'S fluid for 15 to 30 minutes, and then transferring immediately to MERKEL'S fluid (WHITMAN'S modification, with equal parts 1% chromic acid and 0.25% platinic chloride), where they are allowed to remain for one to three hours. They are then washed in distilled water and transferred to alcohol as usual. The treatment with MERKEL'S fluid prevents excessive blackening; the fixation is apparently very faithful and the cilia are well preserved. The sections were mostly 7.5 μ in thickness. The reduced osmium may be still further removed from the sections by leaving the slides over night in a mixture of alcohol and hydrogen peroxide (usually composed of about one part commercial H_2O_2 to three parts strong alcohol, mixed immediately before using). The tissues are by this treatment almost entirely freed of the osmium and left in a suitable condition for staining. The most satisfactory stain was found to be HEIDENHAIN'S iron haematoxylin, with a counter-stain of erythrosin. For study with oil-immersion objectives, the best mounting medium proved to be thickened cedar-oil (immersion oil). This medium hardens slowly, but has the advantage of producing a perfectly homogeneous and transparent lacquer of great permanence and of almost the same refractive index as that of the oil in which the lens is immersed. Very clear and sharp images are thus obtained with the highest powers.

The larvae were fixed by a great variety of methods, of which only a few proved satisfactory. Treatment with 1% osmic acid for 2 to 5 minutes, followed by MERKEL'S fluid for 1—3 hours, gave fair preparations in many instances. The best fixative for general purposes proved to be HERMANN'S fluid (2—5 minutes), followed by MERKEL'S fluid (1—3 hours). In most instances, therefore, the larvae were fixed by the HERMANN-MERKEL method, and the sections (mostly 5 μ in thickness) were stained chiefly with iron haematoxylin and erythrosin as above described, and mounted in cedar-oil. Very clear and sharply differentiated preparations were thus obtained. The tissues were remarkably well preserved, delicate structures

such as cilia retaining, in favorable preparations, almost the norm appearance.

Young larvae of the swarming stage were best prepared b fixation in HERMANN'S fluid for two minutes, followed by MERKEL'S flu for one hour. It is necessary to imbed such larvae as soon as possibl after fixation; otherwise the yolk becomes so brittle as to render practically impossible to secure unbroken sections. If imbedded an cut immediately after fixation, no such difficulty is experienced.

The succeeding descriptions are based upon the examination a very large number of preparations, and all possibility of decep tion arising from imperfect preparations, abnormalities of structure i individual larvae, or failure to observe critical stages, has, it hoped, been avoided.

3. General account of the larval Development with Especia Reference to Segmental Structures.

Arenicola cristata, Stimpson, the species upon which the enti following study has been made, is found in several localities in th neighborhood of Wood's Hole — most abundantly at North Fa mouth, where the extensive sand-flats furnish ideal conditions f its development. The animals are found burrowing at a depth of to 18 inches in the sand, which is deeply saturated with decayin organic matter; and in the summer months are in nearly all in stances well developed and of large size (6—12 inches in length For the larval stages, it has proved necessary to rely upon materia reared in the laboratory, for it has been found impossible in an other way to obtain the requisite quantity of young larvae. I hav frequently searched the sand with a lens in the hope of findin young Arenicolae, but always without success; the larvae, howeve lend themselves so readily to artificial rearing that no difficulty h been experienced in securing in this manner an abundance of spe mens of all stages.

The general characteristics of the adult *A. cristata* are sufficient known from the descriptions of STIMPSON, VERRILL, GAMBLE & AS WORTH, and others. The young free-swimming larvae have be briefly described by WILSON (Johns Hopkins Studies, 1883) and CHU (Z. Bull. Vol. 1, 1897). In the present paper I shall give merely short account of the general characteristics of the larval stage omitting details that do not immediately bear on the problems und

consideration. CHILD (1900) has given an accurate account of the early development within the egg-membrane. My own descriptions will have reference to the metamorphosis and larval development up to a period at which most of the definitive external characteristics are complete.

Very little care is needed in order to preserve the larvae for weeks or even months in a healthy condition, and capable of growth and development. The egg-strings are placed in clean sea-water in large, flat, well-lighted dishes covered by sheets of glass to prevent excess of evaporation, and containing a few pieces of *Ulva* for aëration. It is advisable to change the sea-water at intervals of a week or ten days; otherwise the dishes require very little attention. The organic debris present, derived from minute Algae, particles of decaying *Ulva*, and the bodies of dead larvae, seem to furnish the larvae with sufficient food for the development of a fairly large proportion of their number. Under these conditions, larvae have been kept in the laboratory for periods of 14 to 15 weeks, apparently in a perfectly healthy condition and exhibiting all the normal activities. Development seems, however, to progress more slowly than under the natural conditions, for the largest specimens reared never exceeded a length of 15 mm., although exhibiting in other respects an almost perfect agreement with the adult in appearance, anatomical structure, and behavior. The food-supply is in all probability insufficient for rapid growth; this is indicated by the fact that if kept in sea-water to which carmine powder has been added the larvae usually exhibit a greatly increased rate of growth, especially in the early stages. It would no doubt be possible, by the employment of suitable methods of feeding, to rear them to more advanced stages than the above.

The larvae leave the egg-strings in from two to three days after oviposition, in the form of slightly elongated maggot-like free-swimming organisms (about 0.3 mm. in length), which exhibit a most pronounced positive phototaxis combined with negative geotaxis. As a result of these tendencies they swim rapidly to the light side of the dish and there gather in enormous numbers at the surface of the water. At this stage a larva possesses in addition to the peristomium (which is without setae), three setigerous trunk somites, in each of which are two paired sets of setae corresponding to the notopodial and neuropodial setae of the adult. The notopodial setae of each side are generally two in number, spoon-shaped and spear-

shaped respectively, and are elongated and already capable of a very considerable degree of protrusion and retraction. The neuropodial setae consist of a single short hook on each side of the ventral surface of each setigerous somite. As growth proceeds, additional hooked setae appear ventrally to those already laid down; at the swarming stage, however, a single hook only is as a rule present on either side in each somite. In the third setigerous somite, the setae in the majority of larvae are as yet incompletely formed; as elongation proceeds, however, they enlarge and acquire the same characteristics as those of the preceding somite. Locomotion in the swarming stage is effected by means of the two ciliary rings, prototroch and paratroch; these are connected by a median ventral band of shorter and stouter cilia. The continued activity of the cilia propels the larvae at a uniform rate in a forward direction toward the source of light. On the prostomium are two simple "eye-spots", each of which consists of a compact clump of pigment or excretory granules on the surface of the brain. Large clear cells, apparently of a glandular nature, occupy a large portion of the prostomium, and similar but smaller cells are found at the posterior extremity of the body. These cells probably furnish, in part at least, the glutinous material by the aid of which the larvae later form the rough tubes in which they live subsequently to the termination of the swarming period. In fixed and stained preparations, the cells have a clear and reticulate appearance (Figs. 7 and 8, Plate 22); they diminish and disappear shortly after the beginning of the crawling and burrowing period.

The swarming stage apparently serves simply for the dissemination of the individuals of the new generation, and differentiation of internal organs remains very incomplete until after its termination. The intestine at this period still consists largely of yolk, and its lumen is as yet very incompletely formed. There is, however, a partial subdivision into oesophagus, stomach and rectum — a subdivision which becomes well defined shortly after the beginning of the crawling stage, and persists through part of the larval period and into adult life. Mouth and anus have not yet appeared; the anterior part of the intestine, however, which later forms an eversible proboscis, exhibits already a relatively high degree of differentiation (Fig. 7) — a peculiarity in evident correspondence with the early appearance of its activity and the important part which it plays in larval life. The brain, oesophageal ring and ventral

nerve cord, are continuous with one another and are well defined in the ectoderm of the anterior region of the body. Posteriorly, the ventral cord is less distinctly delimited from the rest of the ectoderm, passing gradually into the undifferentiated tissue of the growing zone. Numerous longitudinal muscle-fibres are closely applied to the inner surface of the body-wall, and fibres derived from this layer are already attached to the inner ends of the seta-sacs. No circular fibres, however, have as yet appeared; these arise later in development and apparently in an entirely different manner from the longitudinal fibrils, i. e. they appear in the ectoderm while the longitudinal fibres are somatopleuric (see below).

The free-swimming stage lasts for a period whose exact duration, in the case of an individual larva, is difficult of determination, but is probably from one to two days at normal summer temperature. At its close the larvae undergo the change which has been described as a metamorphosis; they lose their cilia, sink to the bottom, and adopt a crawling and burrowing and partially tubiculous mode of life which lasts for the remainder of their existence. This change takes place at about the time of formation of the fourth setigerous somite. The larva represented in longitudinal section in figure 8 (Plate 22), is undergoing the transformation; it has already lost most of its cilia; the mouth opening has broken through, and the proctodaeum is in process of formation; the intestinal yolk is evidently undergoing absorption, and the lumen has become more distinct.

The intestinal lumen quickly becomes spacious and continuous from end to end of the body; and in larvae of 6 or 7 somites the yolk has largely disappeared, and the intestine has begun its functional activity. The proboscis, by its continual eversion and retraction, fills the intestine with the surrounding débris, playing at the same time an important part as a burrowing organ, in which form of activity it is assisted by the action of the setae and the muscles of the body-wall. From now on, the habits remain uniform and seem in all essential respects identical with those of the adult *Arenicola*. The material introduced by the proboscis into the intestine is passed through the latter, and the contained organic matter furnishes the sole food-supply of the developing larvae.

With the assumption of the definitive life-habits is associated a change in the reactions towards light and towards the contact of solid bodies. The larvae become negatively heliotropic, exhibiting

a marked tendency to become oriented with the anterior end directed away from the source of illumination, and thus eventually, as a result of the continued crawling movements, to gather in regions remote from the source of light. The other marked peculiarity which now first makes its appearance is a tendency to bring as large a portion as possible of the body-surface into contact with solid bodies (positive stereotropism or thigmotaxis). Larvae freed from adhering solid particles, and placed in a vessel containing loose finely divided débris of any kind, invariably exhibit restless squirming and crawling movements, which continue until the body is once more in contact on all sides with solid particles. When these particles consist of loose sand-grains, diatom-shells or similar light, finely-divided material, the result is that the animal soon becomes enclosed in a rough tube formed by the adhesion of such particles to one another and to the body-surface by means of the abundant secretion furnished by the ectodermal glands. In this manner are formed the rough tubes in which the remainder of the larval period is passed. When the larva is freed from such a tube, the restless squirming movements are resumed, and continue until another tube has been formed. Tendencies such as these, when combined with the characteristic negative heliotropism, will, under natural conditions, plainly favor burrowing movements of the kind observed. Both forms of reaction apparently persist until adult life, and through them the characteristic behavior of the animal is largely determined.

After the close of the swarming period and the assumption of the burrowing habit of life, growth and elongation proceed steadily and uniformly until the full number of somites have been formed. Elongation is due, as in other Annelida, to the activity of a posterior growing zone, which occupies a position immediately anterior to the original situationof the paratroch (Figs. 7—10, Plates 22 and 23 29—33, Plates 23 and 24). The large clear mesodermal and ectodermal cells of this region contain large nucleolated nuclei, frequently found in process of mitosis, and continually give rise anteriorly to new cells which constitute the basis of the newly forming somites. As each somite appears, it is divided off from the one next succeeding by a mesodermal septum; the growing zone thus retains unaltered its distinctive characters, and occupies a constant position relatively to the posterior end.

The short region posterior to the growing zone (post-mesodermal

region or pygidium) retains its characteristic peculiarities in an almost unaltered form throughout the whole of the development (Figs. 7—10, Plates 22 and 23; 29—33, Plates 23 and 24). Its ectodermal cells are small and vacuolated, and contain a granular yellow or brownish pigment, evidently of an excretory nature, which seems to accumulate in this region as the animal grows older — possibly because of its isolation from the rest of the larval body, and its lack of vascular supply. A sharp line of demarcation exists between the ectoderm of this region and that of the growing zone immediately anterior to it. The portion of the intestine contained in this region corresponds to the proctodaeum, and is separated by a valve from the entodermal portion immediately in front (Fig. 33, Plate 24). In its ventral wall is inserted a bunch of strong stiff cilia.

While the posterior portion of the larval body is undergoing elongation in this manner, differentiation of the more anterior somites is in progress. In young larvae (with fewer than nineteen somites), each somite possesses at its earliest appearance — in addition to its section of the intestine and the newly formed muscle-fibres of the body-wall — the rudiments of dorsal and ventral setae; and is separated from its neighbors by complete mesoblastic septal partitions (Figs. 9, 10; 29, 30, 31, 32). As growth proceeds, the septa become incomplete dorsally, and the adjacent body-cavities become continuous with one another. At the same time the setae become fully formed and functional; and the supra- and sub-intestinal blood-vessels, originally simple spaces between the opposed and thinned-out walls of the early mesoblastic somites (Figs. 8, 9), become well defined and of uniform diameter throughout.

The stomach, as described above, is at an early stage (see Figs. 7, 8, Plate 22) already sharply distinct from the oesophagus. Throughout the period now under consideration (from 8 to 18 somites), the oesophagus undergoes a continual backward prolongation in a manner presently to be described, and its region of junction with the stomach is thus gradually shifted backward from the second (Fig. 8) to the seventh somite, its definitive position. In histological structure, the walls of the oesophagus consist chiefly of a single layer of densely ciliated cubical epithelial cells. The stomach is of greater diameter than the oesophagus, and its walls, which are much thicker ventrally than dorsally, are composed of very characteristic large uniformly stained cells of homogeneous appearance. Along its ventral wall extends a ciliated groove continuous anteriorly with

the ciliated cells of the oesophagus and posteriorly with those o the rectum. Posteriorly, the stomach merges gradually in earlie stages into the undifferentiated posterior portion of the intestine later its posterior limit becomes defined, though never with absolut precision; this limit is in the tenth or eleventh somite, where th stomach passes without abrupt transition into the posterior narrowe portion of the intestine. This latter region (rectum) is in its anterio portion completely lined by ciliated cells continuous with those o the stomach; further backward the cilia become restricted to longitudinal groove along the ventral wall; and as the growin region is neared, the cilia finally disappear, and the wall become composed of large clear cells few in number in cross section, an containing large nuclei (Figs. 21, 24, 30, Plate 23).

Both stomach and oesophagus are suspended to the dorsal body wall by a mesentery in which runs the supra-intestinal blood-vessel The sub-intestinal vessel runs along the ventral surface of the inte stine, and presents a characteristic appearance from the larg granular chloragogen cells which invest it. Posteriorly these cell merge gradually into the undifferentiated mesoderm cells of th growing zone (see Figs. 9, 10, 16, 17 etc., 30, Plates 22 and 23).

For the greater part of the above period the septa form in complete transverse partitions extending obliquely backward fron the ventro-lateral regions of the corresponding somites to the sub intestinal blood-vessel. Each septum from the third to the tentl inclusive differs from the rest in being associated with the develop ment of a pair of nephridia, which arise on either side in th mesoblast at the time of formation of the septum, and in continuity with the posterior face of the latter. Each nephridium whicl thus arises soon acquires the form of a minute intracellular tubul in the anterior region of the somite, opening through the septur into the body-cavity beyond by an aperture through which cili project. The two earliest formed pronephridia, those of somites I and V, persist for a short time only, and degenerate at a compara tively early period in the development. The remaining six pair (in somites VI to XI inclusive) are directly transformed into the de finitive adult nephridia. The somites behind the eleventh neve give rise to such pronephridia, so far as my observation has ex tended. It is possible that pronephridia may occasionally appear i these somites, as might be expected from the fact that certain specie exist (e. g. *A. ecaudata*) in which nephridia are normally presen

these somites; but of the numerous larvae that I have examined, ot one has presented this condition, so that its occurrence is at ny rate exceptional. The pronephridia of somites IV and V, on the ontrary, appear with perfect constancy as normal features of the arly development.

In other respects the development of the post-nephridial somites akes place (until the nineteenth somite is reached) in exactly the ame manner as that of the more anterior somites as already escribed. The result is the formation of a chaetigerous anterior ody-region which includes the first eighteen somites. The primi- ve septa are, as above explained, dorsally incomplete, so that the ody-cavities communicate freely with one another throughout the reater part of this region. By the time of attainment of the nineteenth omite it is found, however, that the four most anterior somites ave become separated from the others by muscular partitions which ave arisen at the posterior boundaries of the first, third and fourth omites. These structures are the fore-runners of the three iaphragms of the adult; they do not represent the primitive epta of this region, but are formed independently of these in a anner presently to be described. The body-cavity from the poste- ior boundary of somite IV to the anterior boundary of somite XIX, entirely without septa, and remains in this condition throughout fe, forming the spacious undivided coelom characteristic of the renicolidae.

With the appearance of somite XIX, a marked change takes place n the character of the segmentation. The septum between so- ites XVIII and XIX remains complete, forming a membranous partition hich persists throughout life and divides the anterior chaetigerous ody-region from the region formed behind this limit. The somites ow formed are narrower and shorter than those of the anterior egion (see Plate 24, Fig. 33); they are furthermore entirely desti- ute of setae, and are separated from one another by complete epta. At the beginning of the time of formation of this region, s somites are not conspicuously narrower or shorter than those of ie anterior region. As growth proceeds, however, the characteristic ifferences between the two regions become more evident, and are ell marked by the time the total number of somites has reached ie neighborhood of thirty. With a further increase in the number f somites, the dividing line becomes better defined, until finally, in rvae of 2—2.5 mm. length, the distinction between the two regions

is as distinct as in the adult. The posterior achaetous region undergoes relatively little further differentiation, and its somites exhibit throughout life an almost complete resemblance to one another. The addition of new somites continues in the usual manner until the definitive number has been attained, after which the growing zone loses its distinctive characters and becomes indistinguishable as such.

The disappearance of the growing zone and the cessation of somite-formation take place with great regularity at a stage of about 58 somites. The final number of achaetous somites is thus found to be approximately constant; it may be accurately determined by counting the septa in longitudinal sections of specimens in which the growing zone has disappeared. In such larvae the number is found to range from 38 to 40, and appears never to exceed or fall below this. The total number of somites formed in this species (*Arenicola* seems thus approximately constant (from 56 to 58); in adult specimens, however, the number actually found is generally conspicuously smaller than the above; the deficiency is readily accounted for as due to a loss of a portion of the posterior region by autotomy, which is of frequent occurrence in this section of the body.

Although the posterior region increases in absolute size with the growth of the animal, it undergoes little further differentiation and remains throughout life in a primitive uniformly segmented and achaetous condition. The presence of both dorsal and ventral mesenteries, the regular repetition from end to end of complete septa, the simple uniform structure of the intestine and the regular metameric repetition of the circum-intestinal blood-vessels, are other characteristic features of this region all of them of a decidedly primitive kind. All of these features appear early in the development of the region, and persist throughout life.

With the anterior chaetigerous region the case is very different and very extensive further alterations are necessary before the definitive structure is finally attained. Fig. 30, Plate 23 represents a sagittal section of a larva at a stage of about 8 somites. The metamerism, it will be observed, is expressed in a relatively simple manner: the ventral ectoderm is thickened at segmental intervals, and the septa, all of which are incomplete dorsally (except those just forming), extend from a region a little in front of the constriction upwards and obliquely backwards to the sub-intestinal blood-vessel. The first septum is inserted into the oesophagus immediately behind the proboscis, and into the body-wall almost a

the plane of insertion of the retractor muscles of the proboscis. The oesophagus extends backward through the second somite and joins the stomach near the anterior limit of the third somite; it is evidently undergoing elongation, as may be seen by comparing its extension at this stage with that found in the earlier stages represented in Figures 7 and 8. This backward prolongation is further indicated in the fact that the septa, which in earlier stages occupied planes approximately perpendicular to the long axis (Plates 22 and 23, Figs. 7, 8, 10), have acquired a pronounced slant upwards and backwards. It is to be carefully noted that at this stage the second septum is inserted almost at the anterior boundary of the stomach, while the third and fourth septa, whose line of insertion into the body-wall corresponds with that of the later formed diaphragms, are attached to the ventral blood-vessel at points far behind the anterior boundary of the stomach.

As the oesophagus elongates, the stomach is, as it were, pushed backwards, and the above mentioned backward slope of the septa becomes more marked (Plate 24, Fig. 36). This backward growth has the further result that when the second and third diaphragms are formed (in a manner shortly to be described), each becomes inserted into the oesophagus at a position well in advance of the stomach, although as just shown, this organ originally extends far in advance of the region finally occupied by the oesophagus. The stomach seems during these changes of position to undergo an extensive histolytic disintegration and reorganization to form the tissue of the oesophagus — a fact which must also be taken into consideration in accounting for the changes in the position of septa and diaphragms. These histolytic changes will shortly be described.

The most anterior septum occupies from the first a position immediately behind the line of insertion of the proboscidial retractor muscles (Plate 24, Fig. 35). As growth proceeds, the relations of these two originally independent structures become closer, and at the stage represented in Fig. 36 — in which twelve somites have been laid down — the two have become intimately associated with one another, although still distinguishable on close examination. This association of proboscidial muscles and first septum becomes in later stages more complete, and both eventually enter into the formation of the muscular first diaphragm. The posterior peritoneal wall of the adult diaphragm may be regarded as representing the original first septum.

The ventral portion of the second septum, which, like all t other septa behind the first, is intimately related to the subintestin vessel, carries blood-vessels[1] (which are already laid down in t larva of Plate 24, Fig. 36) from the subintestinal vessel to the vent body-wall of the second somite. These blood-vessels mark through life the original position of the second septum; this structure h thus nothing to do with the formation of the second diaphragm.

The second and third diaphragms, which in the adult form t posterior boundaries of somites 3 and 4 respectively, are form independantly of the primitive septa, and represent entirely dif rent structures from these — as shown by their method of forn tion, histological composition, and relations to internal organs. T second diaphragm begins its appearance at the time of formati of the eleventh or twelfth somite; and in most instances comple its extension across the body cavity during the time of formati of the next two somites. Plate 24, Fig. 36 represents the appearar in sagittal section of a larva of 13 somites, showing the appearar of the newly formed diaphragm under a low power. It will observed that its line of insertion into the ventral body-wall co cides with that of the primitive septum. The latter structure persi for some time after the diaphragm is formed, and can be seen its original position stretching diagonally upwards and backwar to the subintestinal vessel at the anterior border of the stoma The diaphragm on the other hand, extends in a perfectly transver direction completely across the body-cavity and is inserted into t oesophagus in a position a short distance in front of the stomach

The formation of the diaphragm takes place through the inte action of several different and partly independent processes. At i first appearance (in a larva of 12 somites), it has the form of purely membranous partition, inserted into the ventro-lateral regi of the body-wall on a level with the primitive septum, and exte ding transversely to the oesophagus which it joins at a regi

[1] Blood-vessels of essentially similar relations are later laid down in t post-diaphragmal septa; and, as will be shown later, the distribution of segmental blood-vessels in the adult and especially their peculiar relations the nephrostomata, are explained through the fact of their formation in t primitive septa, where they appear as spaces between the two lamellae (Plate Fig. 9; Plate 24, Figs. 32, 33). Since the nephridia are from the first in ir mate connection with the septa, the position of the nephrostomial blood-vess thus becomes readily intelligible. These relations will be more fully dealt w in the division treating of the development of the nephridia.

immediately anterior to the stomach. At this stage, the diaphragm is incomplete dorsally and its muscular fibres are as yet altogether uninformed. Large cells resembling in appearance the mesenchyme cells of earlier stages, are applied to its lateral regions and in the neighborhood of its junction with the oesophagus. These cells evidently play an important part in the formation of the diaphragmal muscle fibres, since the latter make their earliest appearance in regions closely corresponding with the distribution of the cells. At a stage of 12—13 somites, the diaphragm has increased considerably in extent; and muscle-fibres, which from the first show a continuity with the longitudinal muscles of the body-wall, are being laid down in its lateral regions. The extension across the body-cavity has in most larvae become complete by the time of appearance of somite XIV; and an uninterrupted partition, partly muscular, partly membranous, is thus formed at the posterior boundary of the third somite.

At this stage, the essential characteristics of the diaphragm are briefly as follows: at its lateral insertion it is entered on either side by muscle-fibres, continuous with the longitudinal fibrils of the body-wall; these extend completely across the diaphragm, passing below the oesophagus and forming a broad band of muscular tissue extending directly across the body-cavity from side to side. Large nuclei, belonging to the cells above mentioned, are applied to the diaphragm in the region of the muscle-fibres; at the insertion into the oesophagus this organ is constricted, and the muscular fibres and accompanying cells are especially well developed. In its median dorsal and ventral regions, however, the diaphragm still remains membranous and non-muscular.

It seems probable, from the appearances presented at this and later stages, that the cells applied to the originally membranous diaphragm contribute to the formation of its muscle-fibres, i. e., act as myoblast cells, resembling in this respect, the early mesoblast cells of the growing region which give rise to the longitudinal muscles of the body-wall (see below). At the same time the direct continuity of the diaphragmal muscles with those of the body-wall seems to indicate that fibres from the latter are being directly prolonged into the diaphragm by a process of ingrowth. The diaphragmal muscles would on this view have two sources; if, however, the muscle-fibres of the body-wall also increase in number by the activity of myoblast cells (as seems probable), these two modes of origin would be fundamentally identical in nature.

The third diaphragm does not appear until a somewhat lat stage — typically at the time of formation of the sixteenth seventeenth somite. Its manner of formation is in all respects simil to that of the second diaphragm. Its ventral insertion coincid with that of the fourth primitive septum; and the latter, as in t case of the third septum, may persist in its original position f some time after the diaphragm has become complete.

At this stage (17 somites), the oesophageal insertion of the thi diaphragm is immediately anterior to the stomach. The oesophag has thus evidently undergone an elongation of about the length one somite since the time of formation of the second diaphrag In later stages a section of considerable length intervenes betwe the oesophageal insertion of the third diaphragm and the anteri boundary of the stomach. Evidently therefore there is in progre a backward prolongation of the oesophagus. The manner in whi this prolongation takes place is peculiar: it takes place in pa probably by a process of simple backward growth; but it seems be chiefly due to an extensive histolytic transformation of t anterior region of the primitive stomach. The nature of tl transformation may be determined in part from the examination longitudinal sections of larvae of slightly later stages. In a lar of 23 somites, both diaphragms are completely formed; and in t fifth somite, a short distance behind the third diaphragm, the chang alluded to are apparently in active progress. The oesophagus here almost occluded by what appears to be a strong constricti muscle of deeply staining circular fibres, situated at the point junction of oesophagus and stomach. The anterior stomach-regi seems to be undergoing a process of dissolution preparatory to i transformation into oesophageal tissue. The typical unaltered stoma cells are lightly staining, and homogeneous in appearance; th are non-ciliated — except in a groove along the median venti wall (Plate 23, Figs. 16, 17, 30) — and contain a few deeply staini granules. As the region of constriction is neared, however, th appearance undergoes a marked alteration; they become great narrowed and elongated, and drawn out at their inner ends apparen into long tapering flagella. Still farther forward the cells a vacuolated and largely broken down. On passing still farther fo ward the vacuolation becomes less marked; and finally the vacu lated cells pass by a gradual transition into the ordinary cubi ciliated cells of the oesophagus.

Everything thus points to the conclusion that a disintegration and subsequent reorganization of the anterior stomach cells are in progress. In earlier stages, the region of transformation is found farther and farther forwards, and always at the junction of oesophagus and stomach. It thus moves gradually backwards as development proceeds, and the process of transformation apparently ceases only when the definitive limits of the two regions have become finally established. In later larval stages, and in the adult, the oesophagus joins the stomach near the line of separation of the sixth and seventh somites, immediately behind the oesophageal pouches. These latter organs therefore do not arise until oesophagus and stomach have attained their definitive limits.

Some space has been devoted to a description of the above histolytic changes in order to make clear the manner in which the final relations of the diaphragms become established. At the body-wall, the line of insertion of each diaphragm corresponds to that of the original septum to which it corresponds (1st, 3rd and 4th). At the intestinal insertion however, no such correspondence can exist. The whole anterior region of the alimentary canal has undergone a complete alteration from its earlier condition; and the sections of oesophagus bounded by successive diaphragms can therefore in no way correspond to the original segmental divisions. In *Arenicola* the alimentary canal fails indeed to exhibit metamerism in its anterior portions, becoming differentiated as a whole, without reference to segmental limits. In the body-wall, on the other hand, the metamerism — as indicated by the position of the setae, external body-rings, branchiae and nephropores — corresponds closely with that originally laid down in early development. These facts have an important bearing on the question of the possibility of rigorously marking off segmental limits in the adult stages of metameric animals. This can be done only in the case of certain structures in *Arenicola*: the anterior part of the intestine shows no indications of metamerism.

It will be sufficient for the purpose of this paper to give merely a brief outline of the remainder of the larval development, since its details are of interest in the present paper only in so far as they concern the characteristics of the nephridia which will shortly be dealt with at length. As growth proceeds, the division between chaetigerous and achaetous body-sections becomes well-defined, and with the completion of the diaphragms and the definite establishment

of the limits of the intestinal regions, the main characteristics of the adult body are complete. The activity of the growing zone ceases when the larva has attained a length of from 2 to 2.5 mm at which stage the outward appearance and internal organization are essentially identical with those of the adult — further development consisting chiefly in an increase in size and in the completion of histological differentiation. In such larvae, the distribution of the blood-vessels is substantially as in the adult, and the two "hearts" connecting the gastric and subintestinal vessels at the anterior extremity of the stomach, are already differentiated and rhythmically contractile. The gills have appeared in the 7th to 17th setigerous somites (VIII—XVIII) as thin-walled evaginations of the dorso-lateral body-wall containing looped blood-vessels. The otocysts are present on the sides of the prostomium. The six nephridia are visible through the transparent body-wall as sac-like structures in the 5th to the 10th setigerous somites (VI—XI). The dorsal setae and the ventral rows of hooks have the usual characters, but have increased greatly in number as compared with earlier stages. Larvae of 10—15 mm length have undergone little further alteration, except in respect to size and degree of histological differentiation. The gills have become more branched, the "hearts" more conspicuous and better defined, the number of setae has increased, and the external body-rings have appeared. At this stage in fact, the adult organization is essentially complete.

4. Anatomy and histology of the adult nephridium.

The six pairs of nephridia of the adult *Arenicola cristata* are situated in the 5th to the 10th setigerous somites inclusive. These somites correspond however, as seen from a comparison of early larval stages, to body-somites VI to XI, the first somite (peristomium) being without setae, and in the adult, not sharply demarcated externally from the second. Internally, however, its posterior limit is marked by the first diaphragm whose position corresponds closely with that of the original first septum. Between the first and second diaphragms intervenes a region composed of two somites; these are in early stages perfectly distinct from one another; in adult life, however, the bisegmental composition of this region is less evident; it is however, clearly indicated by the presence of two segmental tufts of notopodial setae, as well as by certain peculiarities of the

vascular system (see GAMBLE & ASHWORTH, 1900, Fig. 26). The second diaphragm thus constitutes in reality the posterior boundary of somite III, while the third diaphragm similarly delimits somite IV. This disposition of the diaphragms is characteristic of all the species of *Arenicola* that have thus far been examined, and has its origin in *A. cristata* in a manner that has already been sufficiently described above. The first chaetigerous somite is therefore, strictly speaking, the second body-somite and not the third as GAMBLE & ASHWORTH suppose (1900). It is highly improbable that differences in this respect should prevail among the different species of *Arenicola*; at all events in *A. cristata*, study of the larval stages proves beyond a doubt that the composition of the anterior region is as above described.

The position of the first nephridium in *A. cristata* (body somite VI, or 5th chaetigerous somite) agrees with that of the majority of other species hitherto investigated. In *A. marina*, however, the first nephridium occupies somite V (4th chaetigerous), and its nephrostome perforates the third diaphragm and opens into the body-cavity beyond. It is noteworthy, as GAMBLE & ASHWORTH have emphasized, that this nephridium is typically smaller than the others and frequently much reduced or even absent. Since the larvae of *A. cristata* possess pronephridia in both somites V and IV, it is evident that the present condition has been reached through the disappearance of certain more anteriorly situated nephridia; and the process of reduction is apparently still in progress in *A. marina*. The formation of the anterior diaphragms is possibly in part responsible for the disappearance of the nephridia of this region, since the nephridia are thus cut off from the genital region and deprived of one of their most characteristic functions, that of conveying the genital products to the exterior.

In anatomical structure the nephridia of *A. cristata* are very similar to those of *A. marina*. The large conspicuous nephrostome with its fringed dorsal lip transversed by the nephrostomial blood-vessel, the spacious and uncoiled glandular region, and the large contractile terminal vesicle are characters apparently possessed in common by all species of the genus. The nephrostomial blood-vessel, after leaving the posterior margin of the dorsal lip, passes along the body of the nephridium and is associated in body-somites VII to X with the formation of the gonads. These organs have the characteristic form of strands of cellular tissue traversed by the

blood-vessel and formed by a proliferation of the peritoneal cove of the latter, in a manner later to be described. The positio the anterior margin of the nephrostome corresponds to the boun between two successive somites; in early larval life the nephrost actually perforates the septum and opens in the typical ma into the body cavity beyond. In adult life no septa are pre in the nephridial region; these structures are, however, represe by the segmental blood-vessels which pass between the subintes vessel and the lateral longitudinal vessels. These vessels, with w the nephrostomial blood-vessels are directly continuous, are o nally formed in the intersegmental septa, as already described; in this sense each nephrostome may be considered to open throug life into the next anterior somite. The external opening of nephridium is at the posterior extremity of the somite, dorsal a little posterior to the row of ventral hooked setae.

In its natural position, each nephridium is situated immedia within the dorsal insertion of the transverse muscular bands, which it is partially isolated from the general body-cavity. W these muscles are cut and turned back, the nephridia are fr exposed to view (Plate 22, Fig. 1). Each is then seen as an elong tubular organ with brown walls, terminated posteriorly at the l of the setae by the contractile vesicle, and anteriorly by the l and characteristic nephrostome, conspicuous from the bright processes of its dorsal lip. A slip of transverse muscle is attac to the anterior portion of the dorsal lip and in life binds the nepl stome down in such a manner that its opening is directed inwa Figure 3 represents a nephridium seen in its natural position fr above, after the removal of all the transverse musculature, exc a portion of the attached slip.

The nephrostome is of peculiar and complex structure presents certain peculiarities that apparently have not hitherto b observed in the other species of *Arenicola*. Figure 2, which re sents an enlarged view of the first nephridium of the right s shows the appearance of the nephrostome after its attachments h been removed, and its aperture has been freely exposed to v and spread open. It will be observed that a well defined lin division separates the light-colored non-pigmented nephrostome f the deep brown glandular portion of the organ, a division corres ding to a marked difference in the histological character of lining epithelium of the two regions.

The limits of the nephrostome as such are thus sharply defined in the adult. In early stages, on the other hand, no sharp line of demarcation can be drawn. The above difference of structure must not therefore, as some suppose, be held to indicate that the two regions represent genetically distinct structures which have secondarily entered into intimate association. It will be shown beyond that the two portions develop in direct continuity with one another as portions of an originally single embryonic rudiment.

The dorsal lip is bordered by a series of bright red vascular ciliated processes. These are flattened and somewhat leaf-shaped structures, partially subdivided by secondary finger-like processes, and arranged in a single row from end to end of the dorsal lip. The dorsal lip is somewhat the longer of the two, and is usually somewhat folded in the living *Arenicola*, so that the one-rowed arrangement is not obvious on casual examination. When the nephrostome is slightly stretched, however (as in Fig. 2), the processes are readily seen to have the above-described arrangement, with their broad faces opposed like the leaves of a book. At the anterior and posterior limits of the lip they diminish in size and the number of subdivisions becomes less. Each process may be described as palmate in shape; at its somewhat narrowed base it is attached to the border of the dorsal lip; in structure it is essentially a hollow thin-walled flattened vesicle, lined internally by a connective tissue basement membrane and covered by a columnar ciliated epithelium which passes without transition into the epithelium lining the interior of the nephrostome.

The relation of the processes to the nephrostomial vessel is in reality very simple, and may be readily understood by a reference to the textfigure (p. 362), which represents in a somewhat simplified form a cross section of the entire anterior region of the nephridium. The main portion of the vessel (*nst.v.*) runs along the outer margin of the dorsal lip immediately below the insertion of the processes, and its interior communicates directly with that of the processes, whose bright red color is thus due to the contained blood. No distinction, however, can be seen between the wall of the blood-vessel and the internal connective tissue lining or basement-membrane of the processes, the two being directly continuous with one another, and in reality differing only in their relations to the nephrostomial epithelium. This epithelium is intimately related to that part of the blood-vessel which enters the interior of the processes.


p.d.l.
e.d.l.
nst.v.
e.d.l.
aperture
v.v.l.
m.s.
m.v.l.
b.v.
pn.
gl.neph.
b.v.

xternally, however, the blood-vessel is not covered by epithelium, .t merely by the investing peritoneum of the nephridium (textfig.). ıe relations may, therefore be described as follows: the blood-ssel is bordered by a regular series of short flattened palmate anches or folds, which are covered by the ciliated epithelium of ıe nephrostome and together with this form the ciliated processes f the dorsal lip.

The manner in which these peculiar relations are established ecomes clearer on considering the condition in the later larval tages. The nephrostome in larvae of from 2 to 10 mm. in length s composed of cubical ciliated cells of small size; the margin of he dorsal lip is entire, and along the outer surface runs the early ephrostomial blood-vessel which at this stage is of uniform diameter hroughout its length (Plate 25, Figs. 43, 44, 46—49, *nst.b.v.*). The essel retains this position throughout life; in later stages, however, he portion of the nephrostomial epithelium in contact with the lood-vessels seems to undergo a great increase of surface and to become thrown into folds, which eventually exhibit a typically regular arrangement and definite structure and constitute the processes of the dorsal lip. The wall of the blood-vessel preserves its contact with the nephrostomial epithelium and takes part in the folding process, and hence the interior of the processes necessarily remains in continuity with the lumen of the blood-vessel. The exact manner of the transformation has not been observed, and the above provisional account is largely suppositional, since the margin of the epithelium remains entire in larvae of 10—15 mm. length, the largest I have as yet succeeded in rearing. In other respects, however, the nephrostome of this stage has essentially the same characteristics as that of the adult, and the formation of a system of folds in the manner above outlined would complete the resemblance to the adult nephridium. GAMBLE & ASHWORTH (1898) have

Explanation of Text-figure.

Somewhat simplified cross section through the nephrostome and adjoining glandular region of the adult nephridium. Section-blanc passes about midway between anterior and posterior borders of the nephrostome (see Plate 22, Fig. 2). The figure shows the relations of ciliated processes of dorsal lip (*p.d.l*), nephrostomial blood-vessel (*nst.v*), vesicle of ventral lip (*v.v.l*), and glandular region (*gl.neph.*). The relations of the nephrostomial epithelium to the processes and to the vesicle are shown; also those of the membranous wall of the vesicle (*m.v.l.*) to the epithelium (*e.v.l.*) and to the peritoneum (*p.n.*). *b.v*: blood-vessels; *m.s*: muscular slip attached to dorsal lip; *ex.ep*: excretory epithelium.

published drawings of nephridia of young specimens of *A. mari* respectively 30 and 44 mm. in length (Figs. 16, 18), in which t dorsal processes are just beginning their formation in a manner th corresponds substantially with the above description. In *A. crist* it is probable that they arise at a corresponding stage and in t same manner. It is interesting to note that in certain other speci notably *A. Grubii* and *A. ecaudata*, the processes, while evident bearing the same relation to the nephrostomial blood-vessel, a fewer in number and simpler in structure, remaining apparently i a less differentiated condition throughout life (GAMBLE & ASHWOR 1900).

All indications therefore point to the conclusion that the proces represent simply an elaborate and regular system of folds of t nephrostomial epithelium. The correctness of this view is furth indicated by the identity in structure of the ciliated cells of t processes and of the other portions of the nephrostome.

By the formation of the processes of the dorsal lip, the ciliat surface of the nephrostome is enormously extended and its efficien as an organ for removing suspended solid particles from t coelomic fluid is correspondingly increased. The cilia which cov the processes are incessantly and vigorously active (a conditio presumably favored by the abundant vascular supply), and tend t sweep all suspended solid particles into the interior of the nephri dium. In living Arenicolae it is found that the interior of the nephrostome and the interstices between the processes are occupied by a mass of loosely granular substance evidently collected in *thi* way. On microscopical examination the mass is found to consis largely of broken-down cells containing excretory granules simila to those found in the chloragogen cells coating the ventral blood vessel. It appears probable, in fact, that the chloragogen cell undergo a continual process of disintegration and that the soli excretory products thus set free in the coelom are swept into th interior of the nephridia by the nephrostomial cilia and so conveye to the exterior. The broken-down chloragogen cells are possibl replaced by wandering cells or, more specifically, excretophores which are chemotropically attracted to the ventral blood-vessel an while there extract from the blood the excretory matters; the latte being deposited in a solid form in the interior of the cells an eventually set free in the coelom to be removed to the exterior b the nephridia. Urea and other soluble waste-products seem, on th

other hand, to be excreted by the glandular epithelium of the nephridia. Further evidence of the excretory function of the chloragogen cells will be presented later. At present it is sufficient to note that an important part of the excretory function is apparently performed in the above manner; and it seems possible that the increased ciliary surface provided by the vascular nephrostomial processes has proved advantageous in facilitating the removal of solid excretory particles from the coelomic fluid, and has therefore been acquired in relation to this function (compare ROSA, 1903).

The ventral lip of the nephrostome differs from the dorsal lip in being entire and non-vascular, as in apparently all other species of *Arenicola*. It presents a curious and hitherto undescribed modification in the shape of a large, thin-walled, hollow vesicle (Plate 22, Figs. 2, 4) which occupies its median portion and is prolonged at either end into a tubular region of narrower calibre forming the remainder of the margin of the ventral lip. The entire ventral margin between the anterior and posterior limits of the dorsal lip is thus in reality a thin-walled tube, with a central dilatation, constituting the vesicle in question. The greater part of the wall of the vesicle and of its tubular prolongation is formed of a ciliated epithelium continuous with that lining the interior of the nephrostome (see p. 362). In cross section, the epithelium of the ventral lip is found to curve outward as the margin is reached, and to pass without demarcation into the epithelium covering the outer surface of the tubular region. This epithelium extends around the greater part of the circumference of the vesicle and the marginal tubule; it ceases abruptly, however, at a well defined line an the outer surface (Figs. 2, 6) and the remainder of the circumference is formed of a thin, translucent membrane which at its junction with the wall of the nephrostome becomes continuous with the peritoneal covering of the nephridium. Near its line of junction with the nephrostomial wall, the membrane may show a few blood-vessels derived from the general nephridial network, but towards its junction with the epithelium such vessels are generally wanting.

The interior of the vesicle is typically completely shut off from direct communication with the body-cavity. Occasionally, however, a few ova may be seen within; these may have gained access to its interior as oögonia, and have there undergone further growth and maturation. The thin-walled membranous portion may possibly

exhibit occasional interruptions in its continuity which might ad of the passage of minute bodies such as the early oögonia into interior of the vesicle. Leucocytes appear always to be pres there to a greater or less degree. Definite openings howe placing its interior in free communication with the body-cavity, not exist. The cavity in fact appears simply to represent an enlar lymph-space. It is lined by a thin layer of connective tis containing minute blood-vessels and occasional nuclei, and dire continuous with the membranous portion of the wall of the vesi

The remainder of the ventral lip below the tubular margin like all the non-marginal portion of the nephrostomial wall, compo of (1) an internal layer of cubical ciliated epithelium, lining interior of the nephrostome; and (2) an external connective tis layer bearing blood-vessels and continuous with the periton covering the outer surface of the entire organ.

The manner in which the ventral lip acquires its character structure is at present somewhat uncertain. The vesicle and marg tube make their appearance at a relatively late period of deve ment and are not present in larvae of 10—15 mm., the large have so far reared. The probable manner of their formation however, be inferred from a comparison of the adult and la nephrostomes. In larvae of the above dimensions the nephrost is formed of a single layer of cubical ciliated cells covered ext ally by a thin peritoneal layer (Plate 25, Figs. 43, 44, 48 etc.) possesses an entire margin, along whose dorsal portion runs nephrostomial vessel (Figs. 42—49). The ventral lip is shorter *t* the dorsal lip, but in other respects possesses a similar struct (Fig. 47). The formation of a tubular margin had its origin, in probability, primarily in an increase in the extent of the cili surface. In the adult organ the original border of the nephrost is undoubtedly represented by the line of separation between epithelial and membranous portions of the tubular margin (Plate Fig. 6). The membranous portion apparently represents the orig peritoneal layer, which is here no longer (as originally) in immed contact with the outer wall of the nephrostome, but has bec separated from the latter by the formation of a marginal lyr space of very definite form. The manner in which the pres conditions arose was thus very probably somewhat as follows: The epithelium of the ventral lip underwent a process of extens in an outward direction; if the area of the epithelium increa

ore rapidly than that of the associated peritoneum on its outer de, the former would soon exceed the latter in extent and as a sult would tend at first to curve outward and eventually, with the ntinued growth, to become separated from the peritoneal layer. marginal space would thus be formed between the epithelium and e peritoneum which was originally in contact with it. This space ight later acquire the definite characteristics presented by the ubular marginal space of the adult nephrostome. This space indeed, resents many of the characteristics that we should expect to find ere such its mode of formation. In cross section, its epithelial all is much more extensive than its membranous wall; its epithe- um is continuous with that of the interior of the nephrostome; its terior is closed on all sides toward the body-cavity; and it is lined ternally by a thin layer of connective tissue, continuous with that hich is everywhere present between the peritoneal epithelium and he outer ends of the epithelial cells of the nephridium. Whether r not the conditions of its formation were of the mechanical kind bove imagined, there seems little doubt that it represents essentially large and well defined lymph space between the peritoneum and he outer surface of the nephrostomial epithelium.

The body of the nephridium is of a deep brown color; it is spacious and uncoiled, as in all other species of *Arenicola*, and tapers slightly towards its posterior termination where it joins the large contractile terminal vesicle. The brown color of these regions, like that of the external integument, is due to the numerous excretory granules contained in their walls. The walls of the terminal vesicle are thinner than those of the tubular region, and are typically folded in such a manner as to present a somewhat morula-like appearance, as represented in Plate 22, Figs. 1—4. This appearance is an incidental result of the network-like arrangement of the muscle-fibres which occupy the position of the interspaces between the elevations of the surface. Each inter-muscular area thus tends, in the contracted condition of the vesicle, to project beyond the general surface in the manner represented. Both nephridial tube and terminal vesicle are richly supplied with blood-vessels.

Histology.

The lips of the nephrostome and its interior are lined by a single layer of columnar ciliated cells which show but slight differences of structure in the different regions. Over the surface of the

dorsal processes, the epithelium is folded (Text-figure on p. 36 and is composed of cells of a more columnar shape than thos covering the ventral lip, which are more cubical in appearanc (Plate 22, Fig. 6). The difference in shape is, however, inessentia and is due in all probability to the different conditions of pressur and tension to which the cells are exposed in the different regions the epithelial cells of the processes — being apparently subjected to lateral pressure, as indicated by the folding — are for this reason more columnar; while the cells of the smooth unfolded epithelium of the ventral vesicle are apparently subjected to the tangentia tension characteristic of distended vesicles, and are thus forced t assume a more cubical shape. In all essential respects, however the cells are identical in structure. They are large and clear, c approximately the same size in different regions, and do not contai concretions. Each is bordered by numerous strong cilia (each o which has a well-defined deeply staining basal granule), and th protoplasm exhibits in stained preparations the fibrillar structur characteristic of ciliated cells. As GAMBLE & ASHWORTH hav pointed out (1900, p. 515) the cells are much larger than those o the cubical epithelium of the early nephrostomes. The relative si of the two may be seen by comparing Plate 22, Fig. 6 and Plate 2 Fig. 47, which are drawn to the same scale. It will be seen however, that the nuclei of the early nephrostome are of almos exactly the same size as the nuclei of the adult structure, and occupy a correspondingly large portion of the cell. In their ciliation character of the protoplasm, and general shape, however, the cell are closely similar.

The cubical epithelium of the funnel passes over into the excre tory epithelium of the glandular region by a somewhat abrup transition, although a few of the cubical cells immediately adjoining the excretory epithelium proper may contain granules in their interio and in other respects exhibit a partially intermediate structure.

The large excretory cells of the glandular portion of th nephridium form a single layered excretory epithelium lining th interior of the organ and generally thrown into a well marked systen of internal folds. External to the epithelium is a thin connectiv tissue layer bearing numerous blood-vessels; externally the whol organ is covered by the peritoneum.

In structure the cells agree closely with those of *A. ecaudata* as described by GAMBLE & ASHWORTH. Like most cells of a

excretory function they are much vacuolated and contain numerous deeply staining granules. The nucleus is situated in the middle region of the cell, usually somewhat nearer the base than the outer border. It is large, generally rounded in shape but frequently of irregular outline, and contains a chromatin network of the appearance represented in Plate 22, Fig. 5, together with a conspicuous deeply staining nucleolus which in nearly every instance is seen to contain a well-defined vacuole. The vacuolation of the nucleolus is characteristic of the excretory cells, and also, as will be pointed out later, of the large nuclei of the growing zone, and of the glandular cells of the ventral wall of the stomach. The peculiarity seems to be associated with unusual metabolic activity. It is noteworthy that in the excretory cells the position of the nucleus always closely corresponds with that of the zone of deeply staining excretory granules. — The different regions of the cell show very constant and characteristic peculiarities, as may be seen from Fig. 5 which represents a portion of the excretory epithelium seen in cross section. At its base each cell shows typically a denser structure than at the more peripheral portions, and in fixed and stained preparations, seems composed chiefly of a feltwork of prevailingly longitudinal fibrils — an appearance which seems characteristic of ciliated cells. Typically the basal part of the cell is almost free of granules. Towards the middle region where the nucleus is situated, the protoplasm becomes more vacuolated and is in most instances largely filled with minute round granules which stain an intense black with iron-haematoxylin. Besides these deeply staining granules, there are others (which in some cells may preponderate: see Fig. 5) which stain less deeply and seem to have a greater affinity for the acid stains such as erythrosin. The restriction of the deeply staining granules to the middle region of the cell is very constant, though not absolutely so, and gives rise in cross section to the appearance represented in the text-figure (p. 362). The number of granules, however, shows great variability, some cells being almost free of them while others are so deeply laden as to present an almost uniformly black appearance. The extreme outer portion of the cell is usually, but not always, free of black granules, and is composed of a finely granular, much vacuolated protoplasm which frequently contain numerous lightly staining rounded bodies, apparently of the same nature as the erythrophilous granules just described as present in the middle region of the cell. These bodies range in size from

minute granules to large rounded masses of half the diameter o cell; they are of very variable occurrence, at times very num and at other times almost entirely absent. They apparently r sent the final stage of the excretory products immediately b their extrusion into the lumen of the nephridium. Frequently b of precisely similar appearance are found in the lumen immedi without the cell wall and apparently still adhering to the l Such bodies have evidently recently been extruded. The indica are therefore that the black granules do not represent the stage of the excretory products; but that the latter undergo a fu chemical transformation in the outer portion of the cells before t cast out into the lumen.

The outer rounded margin of the cell usually projects int lumen of the nephridium and bears flagella. The numbe flagella is not, however, constantly one to each cell, as somet held; a single cell may frequently bear a tuft composed of se flagella. The projection of the cell-body into the lumen i general less prominent in cells that contain relatively few excr granules.

The terminal vesicle is also lined internally by excretory of the same general character as the above. Between the l of the epithelial cells and the investing connective tissue laye situated a strong network of muscle fibres whose strands d the surface of the vesicle into the hemispherical elevated areas a described. In other respects, the histological structure of the ve is similar to that of the tubular region. The muscular *networ* derived, in a manner to be later described, from the longitud fibres of the body-wall and is continuous with these.

A comparison of the excretory cells of the adult nephri with those of earlier stages shows various points of agreement disagreement. Fig. 6, Plate 22 shows a longitudinal section thr the wall of the nephridium of a larva of 5—6 mm. The cell will be observed, are much shorter and broader than those of adult nephridium and contain fewer granules, which furthermore to show the definiteness of distribution found in adult individ In the general characteristics of the protoplasm, its vacuolation, its greater density at the base of the cell, the larval excretory agree with the adult; and the nuclei are of almost precisely sir size and appearance, even to the presence in each of a conspic nucleolus which almost always contains a well defined vacuole.

³scular supply of the young nephridium is much less developed an in the adult; and the cells of the terminal vesicle are flatter d contain fewer granules, while the muscular layer of its wall is rgely incomplete (Plate 25, Figs. 45, 55).

The number of granules in the excretory cells of larval stages however subject to great variation, as will be seen by a reference Figs. 46—54 which represent cross sections from the first nephri- ım of a single specimen of somewhat smaller size than the ə from which Fig. 7 was taken. In this specimen the nephridial ls contain numerous and characteristic excretory granules; in other ıpects, however, they are essentially similar to those of the first ɜcimen. The projection of the individual cells into the lumen is ɜsent here as in later stages, though it is less marked.

Later the histology of the larval nephridia will be dealt with in ɔre detail.

5. Development of the Nephridia.

The nephridia develop in close relation to the other segmental ·uctures of the body, especially the septa with which they are ɔm the first in direct continuity. The early somites, as in other ınelids, make their appearance as immediate products of the acti- ty of a growing zone, comparable with the teloblastic region of leech oligochaete embryos. It will be necessary therefore, in order to ake clear the earliest relations of the nephridia, to consider in eater detail the characteristics of this zone and of the region of tive differentiation immediately anterior to it.

(a) Growing Zone.

The appearance of the growing zone in longitudinal section is presented for early larvae in Plates 22 and 23, Figs. 7, 8, 9, 10; for mewhat later stages in Plates 23 and 24, Figs. 30—32; and for the ıge of formation of the achaetous body-region in Plate 24, Fig. 33. ; appearance in cross section may be seen from Figs. 14, 21 and , Plate 23. Its characteristics remain almost constant for the whole riod of its activity; after the formation of 56—58 somites it dis- pears and the formation of somites ceases. Throughout the whole riod of somite-formation the zone retains a constant position a ort distance in front of the posterior extremity of the body, imme- ıtely anterior to the post-mesodermal (pygidial) or former para- ɔchal region. Posteriorly where it adjoins the pygidial region its

boundary is sharply defined; anteriorly it passes by a gradual tr sition into the more fully differentiated region in front. The pygi region thus takes no part in the formation of somites, but is can passively backwards as the body elongates and preserves unch ged its peculiar characters throughout the whole development.

The histological structure of the growing zone may be seen a reference to the figures above mentioned. Cross sections sh that the ectoderm of the extreme posterior region of the body the young larvae is greatly thickened, and of almost uniform meter in all portions of its circumference (Figs. 14, 21 and Plate 23). Figure 14 is a section passing through the posterior re of a larva of 5—6 somites. The section is slightly oblique; dors it cuts through the anterior portion of the paratrochal region, and tr trally through the region of commencing differentiation at the poste border of somite V. The ectoderm of the growing zone is seen be composed of a single layer of large cubical cells contai large clear nuclei, each of which usually exhibits a well-def nucleolus. On passing forward from this zone the dorsal and la regions of the ectoderm are seen gradually to become thinner, cell-limits become less sharply defined, the number of nuclei cross section becomes fewer, and the nuclei themselves become sm and more deeply stained, gradually assuming the usual cha teristics of the tissue-nuclei. The ventral ectoderm also beco thinner anteriorly, although to a less extent than the dorsal lateral regions; and the ventral nerve-cord appears as a fibr differentiation of the inner ends of its cells.

The region of large cells with the large clear nuclei thus dently corresponds to the undifferentiated embryonic region of growing zone, from which cells are divided off anteriorly; in o words, to the teloblastic region. Definite individualized teloblas however are not present in Arenicola at this stage. The nur of cells in cross section is not constant and the cells show no position to become arranged in rows, but simply form an und rentiated zone which passes by a gradual transition into the n differentiated ectoderm in advance. Posteriorly the ectodermal c of the growing zone are, as already mentioned, sharply delim from the small, radially arranged, pigmented cells of the adjoin pygidial region (Plate 24, Fig. 32, 33).

The thinning-out which the ectoderm shows on passing forw from the growing zone is in all probability due in large part t

extension of the body-wall by the contained coelomic fluid, whose osmotic pressure must necessarily exceed that of the sea-water (otherwise collapse would result). The extension of the body-wall in the region immediately anterior to the growing zone is certainly not due purely to a process of cell-multiplication or simple growth. This is shown by the fact that the wall becomes thinner and the nuclei more sparsely distributed at the same time as the superficial extent increases, indicating clearly that the material which is being added to the growing zone is undergoing a re-arrangement and extension of a kind similar to that which would result from mechanical stretching. The expansive force which in this manner stretches the newly formed body-wall and thus causes this region to extend both transversely and longitudinally (see Plates 22—24, Figs. 9, 11, 29—32), is in all probability the osmotic pressure of the coelomic fluid. The process may be compared roughly to the formation of a soap-bubble, where the thick layer of fluid added at the rim of the pipe-bowl (corresponding to the growing zone) is thinned-out and extended in both directions by the pressure acting on the walls from within. The analogy need not be pursued in further detail; it is probably accurate to a certain degree, although in the growth of the larval body-wall material is undoubtedly added at other regions than at the growing zone. The analogy fails for example to explain why the ventral body-wall remains thicker than the dorsal and lateral regions. The above peculiarities of the ectoderm at and immediately in front of the growing zone are however rendered partly intelligible by this comparison.

The space between entoderm and ectoderm at the growing zone is filled with a mass of undifferentiated mesoderm; more anteriorly the mesoderm becomes differentiated and shows the usual division into somites. At the extreme posterior narrow region between the entoderm and the ventral ectoderm are found undifferentiated cells of a characteristic embryonic appearance (Plate 23, Figs. 10, 30, 31). These cells are typically larger and clearer than those situated more anteriorly, and are frequently found in process of mitosis; they possess large clear nuclei each of which typically contains a conspicuous vacuolated nucleolus. They undoubtedly correspond to the teloblasts or mesoblastic pole-cells which terminate the mesoderm bands in so many other species. In *Arenicola*, however, their number is not constant, nor are they sharply defined from the mesoblast cells immediately in front. The extreme posterior mesoderm

might indeed be more accurately described as consisting of a c
nuous mass of undifferentiated protoplasm, containing large n
from which smaller nuclei are divided off anteriorly to form
nuclei of the differentiating mesoderm. The terminal cells ca
thus strictly speaking be regarded as possessing an individua
which persists through development. Described objectively the
ditions are simply as follows: the extreme posterior mesobl
region contains large clear nuclei in process of active multiplica
the nuclei that remain in this position retain their embry
character, while those which are given off anteriorly become sm
and more chromatic (to describe simply the change in appear
and form the nuclei of the newly differentiating tissues. Cel
demarcations are in fact difficult or impossible to observe
early mesoderm of *Arenicola*; the term mesenchyme is th
perhaps more accurate in that it does not imply a definite epit
arrangement of the cells — a condition which is never found i
growing zone of this species.

The undifferentiated portion of the mesoderm is of very lin
extent (Plate 22 and 23, Figs. 9, 10, 29—31); almost immediate
front of the teloblastic region the mesenchyme cells become arra
to form the primitive septum bounding the newly forming so
while the cells applied to the body wall in the same region
fine muscle-fibrils in their interior — the precursors of the l
tudinal muscles of the body-wall (Plate 23, Figs. 13, 14, 21,
The limits of the somites are thus defined very early; when
formed they are very short (Plate 22—24, Figs. 9, 10, 29, 32); t
soon elongate, and attain the normal proportions a short dist
in advance of the growing zone.

The ectoderm and mesoderm of the growing zone thus a
in the possession of large undifferentiated cells containing l
nucleolated nuclei, which form the posterior termination of
growing region and represent the teloblastic or undifferenti
embryonic region of the elongating embryo. The fact that the
of this region are indefinite in number and arrangement — diff
markedly in this respect from the sharply individualized telob
of *Clepsine* (WHITMAN, 1887) and *Lumbricus* (WILSON, 1889) — is
in any degree inconsistent with their being described as telobla
Whether the terminal growing region is constituted of one cel
of several cells seems indeed to be immaterial. The distin
characteristic of teloblastic growth is the presence of a term

.differentiated growing region, which retains its embryonic character roughout development and gives rise anteriorly to the differentiated ssues of the developing organism; and on this criterion the develop- ent of both ectodermal and mesodermal structures in *Arenicola* is pically teloblastic. The entoderm on the other hand does not pparently elongate by teloblastic growth and it does not, pro- erly speaking, exhibit a metameric structure in the same sense as o the ectoderm and mesoderm. Its thin-walled posterior portion is requently drawn out at the lines of insertion of the primitive septa as represented in Plates 23 and 24, Figs. 10, 30—33), and in this ıanner partakes of the early metamerism; but this apparent meta- ıerism seems to be purely incidental and imposed upon the organ om without, and therefore not inherent in its structure. It is oteworthy that in other species that show teloblastic growth the ntestine never seems to develop teloblastically. This fact seems o afford support to those authors (such as HATSCHEK, MEYER, BERGH, GOODRICH) who believe that metamerism in annulate ani- mals is primarily mesoblastic, and that the other metameric ystems of organs have become implicated in a purely secondary nanner.

The large mesodermal and ectodermal cells of the growing zone ıre present, as above described, throughout the entire period of omite-formation. The ectodermal cells are evidently the lineal lescendents of the cells described by CHILD (1900) as derived from he division of the "stemcells", — i. e. of the two symmetrical lerivatives of the somatoblast (2^d of the second quartette of ecto- neres), which form the posterior portion of the somatic plate. These cells at the commencement of the concrescence of the somatic plate form "a transverse band of seven cells lying just anterior to he paratroch and extending in the same direction with it. These ells are large and thick and grow rapidly after division. It seems erfectly clear that we have here what is known as the 'growing ip', i. e. that portion of the body just anterior to the paratroch, which continually gives rise to the new segments and leads to the longation of the larva. The final results of concrescence will be first, to unite the ends of the paratroch, and then the ends of the econd row of cells — the growing tip" (p. 628). CHILD did not ollow the concrescence to its completion; but from the position and relations of the large ectodermal cells of the larval growing region there can be no doubt that these are the direct descendents of the

above band of cells (see CHILD, loc. cit., Figs. 112, 113). The e
retain their large size throughout larval development, showing l
change in character from the early larval stages (three somi
until the completion of the definitive number of somites.

The large terminal mesoblast cells of the larva are similarl
be regarded as the descendents of the two teloblasts of the e
embryonic mesoderm (CHILD, loc. cit. p. 638). The embry
mesoblast bands take up a ventro-lateral position between
entoderm and the ventral ectoderm of the posterior region (Fig. 1
At this stage each band is terminated by a single large telo
According to CHILD "visible differentiation in the descende
the mesoblasts is very slow. The cells retain their emb
appearance till a late stage" (p. 639). The original condition i
each mesoblast band is terminated posteriorly by a single telo
cell does not apparently persist long, for in the larval stag
teloblastic region of the mesoblast always contains several
embryonic nuclei, as above described, which exhibit no con
number or arrangement. Their position, however, between
posterior attachment of the intestine and the ventral ectoderm con
ponds accurately with their position in later embryonic stages
described by CHILD. Between the latest stages described by t
author and the earliest I have examined intervenes a period wh
has not yet been studied with reference to the character of
mesoblast. There can however be no doubt that examination wo
disclose a direct continuity between the teloblasts of the la
embryonic stages, and the posterior undifferentiated mesoderm ce
of the larval growing zone.

(b) Formation of the Somites.

The first larval nephridium appears in somite IV and beco
recognizable soon after the formation of the early septum w
separates this somite from the one immediately anterior to it. Yo
larvae that have just terminated the swarming stage (e. g. Plate
Fig. 8) thus generally exhibit this nephridium in the early sta
of its formation in the mesoderm of the posterior region. Since
of the nephridia arise in exactly the same manner, a descriptio
the changes taking place in the posterior mesoderm of such a la
will apply to all of the early nephridial somites. The mesod
of the non-nephridial somites undergoes closely similar changes

the mesoblastic somites of the whole larval body being indeed med in an essentially unvarying manner.

As the larva advances in development the nephridia of the more erior somites increase in size and undergo a more complete erentiation, at the same time as the others arise in the growing ion. Horizontal sections passing through the nephridial region such larvae may in fortunate preparations show the entire series nephridia, from the early undifferentiated rudiments of the region oining the growing zone, to the well-defined nephridia of the re anterior somites. Fig. 29, Plate 23 represents such a prepara- n in the case of a larva of about eleven somites. On the right e the section passes slightly ventrad to the ventral hooked setae I shows the larval nephridia from the first (in somite IV) to the th (somite IX); the undifferentiated rudiment of the 7th (5th defini- e nephridium) is just visible at the angle between the newly ming septum and the bodywall. In Figs. 16 to 22, inclusive late 23), have been represented cross sections of the successive phridial somites and the post-mesodermal region of a larva in ich somite IX is in process of formation. The different stages in e early development of a nephridium can thus be studied to vantage in single larvae of this and similar stages. The develop- ent of the nephridia is however so intimately bound up with that the somites that it will be necessary first of all to consider the ter process in detail.

The first changes, preparatory to somite-formation, that take ce in the posterior undifferentiated mesodermal region consist in multiplication and subsequent rearrangement of the mesodermal ls, which at first are tightly packed together at the posterior angle tween the intestine and the body-wall (Plates 22—24, Figs. 7—10, —33). The manner in which this rearrangement is accomplished y be seen from the above drawings, especially the sagittal tions (Plate 23, Figs. 10 and 31), and the horizontal sections ate 22, Fig. 9 and Plate 24, Fig. 32). The closely packed posterior soderm is here seen to be bounded anteriorly by a roughly defined nsverse partition whose cells are directly continuous with the senchyme cells lining the cavity of the more fully formed adjoin- g (somite VI, Fig. 9; somite VIII, Fig. 10). This partition, therefore, presents the posterior wall of the somite next in front of the one process of formation, and defines the position of the future inter- gmental septum, of which structure it in fact forms the anterior

lamella, as will be shown beyond. As the number of mesench cells behind this partition increases — chiefly by the multiplica of the large polar nuclei — an irregular hollow space app between the cells in the neighborhood of the partition, and gradu enlarges, apparently in part by the absorption of fluid from adjoining coelomic cavities. The space so formed represents early coelomic cavity of the somite which is in process of formati it becomes progressively better defined as growth proceeds, wi the layer of mesoblast forming its walls becomes irregularly appli to the surface of adjoining structures — anteriorly to the poste wall of the next somite in front, externally to the ectoderm of body-wall, internally to the wall of the intestine and posterior the densely packed mesoderm of the growing region behind partition formed by the apposition of these mesoblast cells posterior wall of the somite in front represents the earliest con of the newly forming intersegmental septum, and thus define anterior boundary of the newly forming somite. The early sept thus consists essentially of the opposed walls of this incipient som and of the somite in front. Its characteristics and relations to nephridia will shortly be considered in detail.

The bilaterality of the early mesoderm is not very decid pronounced. A ventral mesentery is apparently not formed in anterior region of the body and the coelomic cavities of the sides are therefore from the first continuous below the intes Dorsally, however (Plate 23, Fig. 13), the right and left coel cavities remain distinct, and the adjoining walls (which soon ass a membranous character) become applied to one another above intestine, forming a bilaminar membrane, whose lamellae are the conditions of their formation necessarily continuous with mesoderm covering the intestinal and body-walls, and with lamellae of the primitive septa. This membrane constitutes dorsal mesentery which persists throughout life and suspends intestine to the dorsal body-wall.

Before considering in greater detail the characteristics of primitive septa, the formation of the dorsal and ventral bl vessels may be briefly described, since these structures appear early and apparently play an important part in the nutrition of growing region. The dorsal blood-vessel is early formed as a s between the opposed lamellae of the dorsal mesentery (Plat Figs. 13, 20 and 21). It does not, however, appear to become

fined so soon as the ventral vessel, being possibly less necessary early stages than the latter, in which the blood flow is from fore backward towards the growing zone. This vessel, as will seen from Fig. 9, Plate 22, early appears as a well defined nnel extending almost to the posterior extremity; it arises as a ace between the intestinal wall and the mesoderm and soon be- nes enclosed on all sides by the extension of the mesoderm cells. walls are from the first continuous with the original septa, a fact significance, since the segmental blood-vessels arise in the septa l in this manner acquire their characteristic relations to the phridia, as will be explained beyond.

Mesoderm cells remain applied to the wall of this vessel, and development proceeds they acquire a voluminous and vacuolated pearance, and pigment or excretory granules become deposited in eir interior. These cells constitute the chloragogen cells; they e thus in reality modified peritoneal cells, which throughout life at the ventral blood-vessel and play an important part in the ocess of excretion (cf. ROSA 1903). They have in most larvae eady assumed their excretory character in the third somite in nt of the growing zone, and from this region they may be seen pass backwards by a gradual transition into the posterior un- fferentiated mesenchyme cells (Fig. 9). Their characteristics will be nsidered in greater detail below.

Dorsal and ventral blood-vessels soon acquire open communica- n with each other in the growing zone, thus completing the circu- ion and providing more fully for the nutrition of this region. e nature of this communication may be seen from Fig. 24, Plate 23, hich is a cross section through the posterior region of a larva of —12 somites, passing through the region of the primitive septum tween somites XI and XII. Dorsal and ventral vessels are here en to be connected by a blood-sinus which completely surrounds e intestine. A sinus of this nature is typically present in larvae this and later stages; it consists of a simple space between the estinal wall and the mesoderm, and is situated usually at the sertion of the most posterior primitive septum, but frequently (see ate 23, Fig. 30 and Plate 24, Fig. 32) at the membrane which separates e posterior mesoderm from the space between proctodaeum and toderm in the post-mesodermal region. The sinus varies greatly its size and in the amount of blood that it contains and at times ems to become obliterated. Apparently it is formed as required

at the zone where least resistance is offered to the passage of b between the two vessels, and thus represents no sharply define persistent channel. The branching blood-sinus which in later envelopes the stomach-region seems to be formed in essentially same manner.

The primitive septa are formed, as above described, by apposition of the adjoining walls of two successive mesobla somites. The walls of these early somites are, however, no formed of a regular epithelial layer of well-defined cells, but to appearance consist of a continuous syncytial layer of protoplasm taining numerous nuclei (Plate 23, Figs. 13, 14, 23, 24), and d applied to the surface of adjoining structures. Each nucleu for purposes of description be regarded as belonging to a cell, but definite cell-walls, at this stage at least, are distinguishable.

The structure of a septum at its earliest distinguishable p may be best seen in Figs. 9 and 10, Plate 22 and 23, and Fig Plate 23. At this period of its formation the septum forms an gular nucleated partition, separating the cavity of the newly-for somite from that of the somite next in advance. At this stag bilaminar structure can with certainty be discerned in it, and constituent cells, which resemble in all respects the other mesod cells of this region, seem to be entirely undifferentiated and si to one another. As growth proceeds, however, the coelomic ca increase in size and in transverse diameter in the manner desc on page 374; the septa are thus subjected to a stretching pr and increase rapidly in superficial extent, becoming at the time thinner and better defined. During this process the op faces of the cells of the two septal layers (which are now u distinguishable from each other) assume a more membranous char becoming at the same time more deeply staining and apparen a denser consistency (Plate 24, Fig. 32). As the septum cont to extend it acquires more and more of the character of a membrane, and the nuclei of its constituent cells are moved fa and farther apart. Finally the septum acquires the appearan a thin, sharply defined, membranous partition extending acros body cavity, and to whose surface the original nuclei (each of w is still surrounded by a small quantity of undifferentiated protop are applied at infrequent intervals (Plates 22—24, Figs. 9, 29-

At the junction of the septum with the body-wall and wit

stinal blood-vessel the two closely applied lamellae are usu-
en to be slightly separated from each other where they pass
ntinuity with the peritoneal membrane. In the larva repre-
in section in Fig. 9 the interlamellar space is continued for
ht distance outward from the lumen of the sub-intestinal
vessel; this space represents in all probability the beginning
segmental blood-vessel which later passes to the body-wall
iverses the dorsal lip of the nephrostome. The septum is seen
s figure (and also in Fig. 10, 29, 31, 33, Plates 23 and 24)
ome continuous at its lateral margin with the lips of the
stome. At its insertion into the body-wall its cells become
ious with those lining the body-cavity.

he line along which the septum is inserted into the body-wall
ally considered to mark the boundary between two successive
s. This line occupies in *Arenicola* a constant and definite
n with reference to the other segmental structures of the
ping larva. During the formation of the mesoblastic somites
ody-wall also undergoes a division into segmental regions.
netameric structure of the ectoderm is most conspicuously
ed by the setae which appear at an early stage in the
ils between the successive septa. The seta-sacs which give
these structures appear laterally in the ectoderm in a position
y behind the middle of the interseptal region (Fig. 32, Plate 24).
entral ectoderm in the neighborhood of the nerve-cord also
exhibits a metameric structure, becoming slightly thinner or
cted at segmental intervals. The septa are inserted ventrally
a transverse line passing immediately in front of the con-
n (Plate 23, Fig. 30). This line, therefore, if the above crite-
f segmental limits is accepted, marks the boundary between
sive somites. Later in development, when the septum has
eared as such, the position of the line of demarcation corres-
closely with that of the nephrostome and its associated branch
segmental blood-vessel — both of which structures, as already
arise in intimate connection with the primitive intersegmental
.

he septum, when first formed, constitutes a complete partition
n the body-cavities of adjacent somites. Soon after its forma-
owever, it becomes incomplete dorsally, as already mentioned
), and for the remainder of its existence it is found only in
ntral portion of the body-cavity as an incomplete membranous

partition, extending from the ventral blood-vessel to the ventro-lateral body-wall. It usually becomes incomplete at the second or third somite in advance of the growing region (Plates 22 and 23, Figs. 8—10, 29, 30), apparently as a result of the thinning and eventual disruption of its dorsal portions. The conditions that lead to this result seem to depend in part upon the progressive increase in the transverse diameter of this region of the body, associated, as above described, with an increase in the extent of the body-wall. A further reference to Plate 23, Figs. 16—21, which represent sections through successive somites of a single larva, will show that the increase in transverse diameter is dependent chiefly upon the extension of the thin-walled dorsal and lateral regions of the body-wall. It seems, therefore, probable, as above suggested, that the septa first become incomplete in the dorsal region largely in consequence of the undue stretching to which they are thus subjected, which has caused here a thinning and eventually a disruption and disappearance of the septa. On the other hand, their ventral portions persist for a very considerable period, and play an important part in later development in connection with the nephridia and nephrostomial blood-vessels.

(c) Formation of the Nephridia.

I shall now proceed to a description of the histogenetic processes by which the nephridia are differentiated out of the originally uniform somatopleuric layer of the mesoblast. In the region immediately adjoining the pole-cells this layer, as described above, is composed of a continuous nucleated protoplasm without cell-boundaries and without visible differentiation; at a slightly more anterior position, however, where the somatopleure passes into continuity with the newly forming septum, the differentiation of the longitudinal muscle-fibres of the body-wall is already in progress. These structures make their first appearance at a very early period, in that portion of the mesoblast which is in immediate contact with the ectoderm of the body-wall (Plate 23, Figs. 13, 14, 21, 24). The fibres first become visible in two ventro-lateral areas on either side of the ventral nerve-cord, which is just beginning its appearance (*n.c.*); and at a slightly later period they appear also in two dorso-lateral areas on either side of the dorsal mesentery. Along the line of insertion of the dorsal setae muscle-fibres do not at first appear.

At the earliest appearance the fibres have in cross section the pearance of a row of deeply-staining dots at the boundary between oderm and mesoderm (Plate 23, Figs. 13, 21, 24). At a slightly er stage (Figs. 15, 23) the dots have become larger and better fined, and have assumed the typical appearance of triangular dge-shaped structures, which stain very deeply in the erythrosine d project into the protoplasm of the mesoblast cells. In the ventro-eral region the mesoblast nuclei have become arranged in a single er, and in cross section the mesoblast has assumed the appearance, ich is very typical for this stage, of a row of nucleated cells, eparated by cell-walls, and containing in their interior deeply ming longitudinal muscle fibrils (Plate 23, Figs. 13, 15, 19, 23). s evident that the longitudinal muscle-fibres are originally laid n in the interior of these cells, which may therefore with pro-ty be termed myoblast cells. The cells, however, it is to be ed, present no constant and regular arrangement and are not rply defined from one another. At this stage the longitudinal scles are thus composed of fibrillae applied to the body-wall and ered internally by a nucleated layer of undifferentiated protoplasm coplasm). The condition recalls that which exists throughout in certain other species, particularly the Nemathelminthes. The nsverse muscles on the other hand are not visible until a much er stage; they appear externally to the longitudinal muscles, and parently in relation to the inner layer of the ectodermal cells[1]. e above myoblasts at all events have nothing to do with their rmation.

Cross sections passing through the region immediately behind e early septum show the above-described appearances in a charac-ristic manner (Figs. 13, 15, 19, 23). Longitudinal sections (Plates 22 and , Figs. 9, 10 and 31), passing through the ventro-lateral regions, show at the mesoblastic layer is thickest and best defined at the junction septum and body-wall. It is in this region that the embryonic diments of the nephridia (*n*) first become distinguishable from the st of the mesoblastic layer. It must be emphasized however, that ere is at first no setting aside of special nephridial cells; nothing mparable to the septal nephroblasts or "funnel-cells" described by

[1] Ed. Meyer (1901) gives a summary of the evidence on this point, from ich it would appear that an ectodermal origin is the rule, if not invariable, the circular body-musculature of annelids.

VEJDOVSKY, WILSON and BERGH and more recently by I (1902). The nephridia simply become gradually differentiat of an originally uniform and homogenous mesoblast which in stages presents no definite cell-boundaries. The differentiati the nephridium starts from the angle between septum and wall; but at first no visible differences exist between the ne blastic mesoderm and the adjoining myoblastic or septal meso

The first signs of the appearance of the nephridium are v usually at the second septum in advance of the growing Young larvae of five somites thus as a rule exhibit the first h nephridium in the earliest stages of its formation. The longit section represented in Fig. 9, Plate 22 shows the differe 2d nephridium in somite V. At this stage the rudimenta has the form of a somewhat triangular mass of protopla tinuous anteriorly with the fourth septum. Fig. 10, Plate 23 a similar stage in the formation of the fourth nephridium; and F similarly represents the rudiment of the sixth. The continuity the septum at the junction of the latter with the body-wall is haps the most characteristic feature of the early nephridial rudi

The histological differentiation of the nephridial cells c studied to better advantage in transverse sections. In Pla Fig. 13, which represents a section through the fourth septal of a larva of 5—6 somites, the somatopleuric cells of the v lateral region are entirely alike, with nothing to distinguis future nephridial region from the other portions of the mes The protoplasm is homogenous and non-granular, the nucl similar to one another in size and appearance, and cell-wal indistinguishable. At this stage in fact no essential difference between the cells of this intersegmental region, and those intersegmental region of somites II and III, where no nephrid formed (Plate 23, Fig. 12). In Fig. 15, which represents a through the beginning of somite VI in a larva of about eight s one of the cells of the right side is distinguished from the by the presence of a small space, bordered by deeply st protoplasm, which has made its appearance in the immediat nity of the nucleus. This minute space is the first indication appearance of the nephridial lumen, which, it is of interest t always seems to arise in the neighborhood of the nucleus. I respects the cell which shows this peculiarity is entirely sim the adjoining purely myoblastic cells, even to the posse

muscle-fibres in its interior. Fig. 23, Plate 23 shows a ilar condition. The nephridium is thus differentiated out of a tion of the original mesoblast which in its early stages possesses same properties and undergoes the same transformations as the er portions of the layer.

At a slightly later stage the nephridial cells become more readily tinguishable from the adjoining purely myoblast cells. The lumen comes more distinct and the body of the differentiating organ increases in size, apparently in part at the expense of the protoplasm the adjoining mesoblast cells, which become visibly smaller. In section the nephridial rudiment is soon observed to be more minent than the adjoining myoblastic cells, and its boundaries ear better defined (Plate 23, Figs. 23 left, 19 left). Its protoplasm becomes vacuolated and minute granules in many cases begin make their appearance in its interior; the lumen becomes progressively wider and better defined and the cilia become visible. The section represented in Fig. 19 (3 somites from the growing end) the nephridium of the left side is somewhat more advanced than that of the right and is becoming sharply defined from the joining myoblast cells, which have perceptibly diminished in volume. The next somite of the same larva (Fig. 18) the boundaries of the nephridium are quite definite, and flattened mesoblast cells are being applied to its surface to form the outer peritoneal covering the organ. The nephridium in the somite next in advance (Fig. 17) is still more highly differentiated; the lumen is more spacious and the excretory granules are more numerous. At this stage the organ has the form of a simple tubule with an intracellular lumen opening anteriorly through the septum into the cavity of the adjoining somite. The progressive histological differentiation of the early nephridia is well shown in the series of sections represented in these drawings (Plate 23, Figs. 16—21).

As the nephridium advances in development the myoblast cells gradually diminish in volume, while at the same time the muscle-fibrils show their interior increase in size (Figs. 16—21). Eventually the undifferentiated protoplasm or sarcoplasm is reduced to a thin layer occupying the interstices between the fibrils and containing here and there a flattened nucleus. In later stages (Plate 23, Figs. 25—28) the muscle fibres become much more numerous and more closely set together, and the sarcoplasm is reduced to an inconspicuous interfibrillar matrix in which the fibres are imbedded. The innermost layer

of the sarcoplasm seems to become membranous and to contribu
the formation of the peritoneal lining of the body-cavity.

The nephridial rudiment can as a rule be distinguished be
the septum has assumed a definite membranous structure (Plate
Fig. 31). It is therefore difficult to decide whether or not
nephridium is derived chiefly from one or other of the two lame
of which the septum is composed, or from both. From all appearan
however, it seems clear that neither lamella is exclusively concer
in its formation but that both contribute in part; and that the p
septal mesoblast also participates in the process. The protopl
of the nephridial rudiment and that of the septal cells are at f
continous with one another; it is only later that the boundar
the two become sharply defined. The fact, however, that
very earliest appearance the nephridial lumen communicate
the body-cavity of the preceding somite proves that the meso
wall of this somite enters, in part at least, into the formatio
the early nephrostome. It cannot, however, be said that the nep
stome is formed exclusively from the posterior wall of this so
The early nephridial rudiment is in fact a continuous and undi
structure without regional delimitations, and it is not until
later that the nephrostome becomes distinct from the glan
portion of the organ. It is impossible, therefore, to say that
definite portion of the nephridium is derived from one somite
another portion from another somite. It is even impossible in
early stages to sharply define the limits of the mesoderm of su
sive somites, since the differentiation of somites is itself a gr
process, and the nephridial rudiment is already laid down b
the segmental limits are sharply defined. It is, however, true
in its early undifferentiated condition the nephridium is conti
with mesodermal tissue which later forms portions of two succ
somites. The organ, however, is differentiated as a whole w
strict reference to segmental limits. Even cell-limits, as al
pointed out, are not sharply defined until the organ has al
acquired many of its most distinctive characters. The whole
early nephridium is, however, strictly mesoblastic and the ect
plays no part whatever in its formation. Later in developme
will be shown, the ectoderm apparently contributes to the forn
of the terminal vesicle; but as originally laid down in early de
ment the entire organ is strictly mesoblastic.

The structure of the early nephridium, as present in the

h somite in advance of the growing zone, is shown in longitu-
section in Fig. 11, Plate 23. The anterior end of the organ
ds across the thin ectodermal region just behind the segmental
and its lips are continuous with the remains of the primitive
n (not shown in Fig. 11). The organ consists of a straight
nched tubule whose walls are composed of a continuous vacuo-
protoplasm containing excretory granules. An axial ciliated
is present, which freely communicates with the body-cavity
or to the septum by means of a minute opening through which
lia project. Nuclei are present in the walls of the organ,
as yet shows no division into well-defined cells. As yet the
ı does not communicate with the exterior, although the posterior
nity of the organ is closely applied to the ectoderm and has
ently become continuous with the latter. Fig. 34, Plate 24
sents a tangential section through the ventro-lateral region
arva of 13—14 somites, showing the two most anterior nephridia.
nephridia are the 2nd and 3rd larval nephridia (2nd provisional
1st definitive, of somites V and VI respectively), the first pair
rval nephridia having already disappeared. The nephridia
increased in size considerably as compared with the stage
ented in Fig. 11, Plate 23, but are otherwise essentially as
bed above. The posterior fusion with the body wall is here
ı; the fusion takes place slightly behind and internally to the
f ventral hooked setae. The lumen is incomplete posteriorly.
iorly the continuity of the nephrostomial border with the septum
arly shown.

At this period the nephridium is, in brief, a simple tubule with
ellular lumen, of the kind frequently described in other larval
lids (cf. MEYER, 1887—88, LANG, 1889). It is peculiar, however,
ing anteriorly in open communication with the body-cavity. In
eculiarity it presents a distinct contrast to the pronephridia de-
d by MEYER and LANG, which are described as closed inter-
r and as bearing a marked resemblance to the flame-cells of
llaria — a resemblance which has formed the basis of much
genetic speculation. The pronephridia of *Arenicola*, however,
ss a distinct coelomic aperture of the kind described, as I have
ved with perfect clearness an indefinite number of times. The
ior region of these organs in fact, together with a portion of
djoining septum, constitutes the primitive nephrostome, from
ı the adult nephrostome is directly derived, as will shortly be shown.

It is noteworthy that at the stage represented in F
Plate 24 the two foremost pair of nephridia are in all respect
although the anterior of the two later on completely dege
and disappears, while the posterior continues its develop
forms the first definitive nephridium of the adult. The firs
nephridium (in somite IV) has in most larvae already di
at this stage. The time of its disappearance is variable; i
larvae it has already disappeared at a stage of 9—10 somi
in others it may persist in a recognizable form until fifteen
somites have been attained. The second larval nephridium,
correspond to the first definitive nephridium of *A. mari*
degenerates and disappears, but not until a considerably la

(d) Later Development of the Nephridium.

The later development of the nephridium consists chiefly
an increase in size and an accompanying division into well-
cells with definite boundaries; (2) in a differentiation of the
region of the nephridium, and apparently also of a portion
adjoining septum, to form the nephrostome, and in the establis
of characteristic relations with the blood-vessels; and (3)
formation of the terminal vesicle. Each of these processes w
considered in order.

Subdivision into Cells.

In nephridia of the stage represented in Fig. 34, Plate 24
definite cell-boundaries are visible, although the protoplasm ha
typical vacuolated structure and contains excretory granules
remains none the less continuous and undivided from end to
of the organ; its nuclei are small and as yet differ only slig
from the neighboring mesodermal and ectodermal nuclei of the b
wall. The lumen retains this intracellular character for a consi
able period. Fig. 25, Plate 23 represents a cross section of the
nephridium (somite XI) of a more advanced larva than tha
Fig. 34; the organ has increased considerably in size and its p
plasm, together with the large nucleolated nuclei, shows the stru
characteristic of fully differentiated and functional excretory
As yet however, cell-limits have not appeared and the lumen ren
intracellular. Fig. 37, Plate 24 represents a longitudinal sectio
the corresponding nephridium (somite XI) of a slightly more adva
larva in which the organ is beginning to show a subdivision

A cross section of a similar nephridium is represented in
', Plate 23. The cells are large and few in number, and
l large nucleolated nuclei. At the narrower posterior region
y nephridia of this stage the lumen may remain intracellular,
though its anterior region may be intercellular and bounded
or three distinct cells in cross section. As the cells increase
ber, the entire lumen becomes intercellular. The distinction
ı the two forms of lumen seems thus purely artificial and a
ous gradation can be traced.

e subdivision into cells takes place apparently in simple rela-
the increased calibre of the lumen of the organ. Accordingly
und, as growth proceeds, that the lumen becomes bordered
ell-defined layer of cubical excretory cells whose number in
ection is in general directly proportional to the measure of
cumference of the lumen. Figs. 27 and 28, Plate 23 show
ive stages in the formation of this epithelium. Fig. 28
nts a cross section through the body of a nephridium of almost
ne stage as that of Fig. 42, Plate 25. Five cells are already
in cross section, and their inner ends are beginning to pro-
to the lumen in the manner characteristic of latter stages.
', 46—54, Plate 25 and Fig. 6a, Plate 22 show still more
d nephridia with a greatly increased number of cells in cross

ce the length of the nephridium in *Arenicola* is limited, never
ng that of the somite to which it belongs, the only way in
t is possible to secure the necessary increase in the extent
excretory surface is by an increase in the calibre of the
Since each nucleus is in direct physiological relation to only
d portion of the excretory protoplasm, the increase in calibre
a corresponding increase in the number of cells in cross
The cells retain an approximately uniform size (as may
comparing the above figures), and increase in number
itly by direct division) as the epithelium increases in extent.
are never seen in the excretory epithelium; and it seems
at the cells when they exceed a certain limit of size, undergo
e subdivision, preceded by an amitotic division of the nucleus.
not directly observed the several stages of this process; the
however, are frequently seen in an irregular or constricted
nd subdivision probably simply represents the extreme of
condition. The formation of a cell wall between the two

nuclei thus formed would complete the process. The structu the larval excretory cells varies but slightly at different stage the adult nephridium the cells have become very numerous greatly compressed in a lateral direction, assuming a marked colu structure; in consequence of this, apparently, the different re of individual cells may present more marked differences from another than are found in the broad cubical cells of the l nephridium. In essential histological character, however, the l and adult cells seem to differ but slightly from each (see p. 370).

Nephrostome and its Associated Blood-Vessels

In the formation of the nephrostome both the anterior p the original nephridium and a portion of the adjoining sept part. From its earliest appearance, as above shown, the nep possesses an anterior opening into the body-cavity of the next s but at first the marginal region bordering this opening is i respect different from the remainder of the organ. When subdi into cells begins the cells of the anterior region at first en resemble the others in structure and appearance; as growth ceeds, however, subdivision progresses more rapidly in these than in those behind and as a result the more anteriorly si cells become smaller and more cubical. At the same time the their distinctively excretory character and the protoplasm be more homogeneous in appearance, while the vacuoles and exc granules disappear (Plates 24 and 25, Figs. 37, 39, 42).

In Figs. 37 and 39, a portion of the adjoining septum a to be assuming the character of a cubical epithelium and t be contributing to the extension of the dorsal lip of the funnel limits between septum and nephridium proper are in fact not s defined in early stages (Plate 24, Fig. 34); and until the f epithelium is definitely formed, it is impossible to say whe two become continuous. Its hinder part however is unquesti formed from the anterior cells of the nephridium proper (Figs. 3 42), while its marginal cells, especially those of the dorsal lip to be formed in the region of the original septum. The clos tion of the dorsal lip to the septal structures, especially the vessels (a relation which persists throughout life), is thus exp The anterior portion of the nephridium becomes in this r composed of an epithelium of small cubical ciliated cells (sh

ss section in Plate 25, Figs. 43 and 44). The cilia of these cells come shorter and more numerous (compare Plate 24, Figs. 38, 39; te 25, Figs. 42, 46—49), while at the same time the nuclei become aller and more rounded, and the nucleoli characteristic of the retory cells usually disappear. Eventually a well-defined funnel ion is formed whose cells are sharply distinct from the excretory s behind (Figs. 46—49).

During the earlier stages of the above transformation the intermental septum undergoes changes which lead to the formation he segmental blood-vessel. The septum at the beginning of the od in question extends obliquely forward from the ventral region he stomach (where it is continuous with the sub-intestinal blood-el) to the ventral body-wall, forming a thin membrane which nds laterally to a point slightly above the level of the early hrostome (Plate 25, Fig. 43). In its lateral portion its border omes continuous with the lip of the early nephrostome, as above cribed — the anterior face of the septum being directly continuous h the epithelium of the funnel — while the posterior lamella ses directly into the peritoneum covering the nephridium. Figs. 37 9, Plate 24 show the relations of septum and nephridium at the ly period of differentiation of the nephrostome before definite od-vessels have appeared.

Each segmental blood-vessel appears in the septum (see Plate 25, . 43, in which portions of the original septum yet remain), origily as a space between the two lamellae of the latter. The vessel ins its formation at the junction with the sub-intestinal blood-sel (Plate 22, Fig. 9) and gradually extends outward to the body-ll where it joins the lateral longitudinal vessel, also formed about s time. (The relations of the segmental vessel to the nephridium in part shown in Plate 25, Figs. 42—44). Near its junction h the body-wall the main vessel gives off a branch (the nephro-nial vessel) which curves back and passes inward and backward ng the dorsal lip of the nephrostome between the funnel-epithe-m and the peritoneum, to the glandular portion of the nephridium ng which it passes as the main nephridial vessel. The nephro-nial vessel is thus directly continuous with the segmental vessel l is formed in essentially the same manner as this latter, repre-ting in reality a space between the posterior septal lamella presented by the peritoneum) and the anterior lamella at the ion where the latter joins the funnel-epithelium with which it is

continuous. As the epithelium increases in extent the vessel acq its definitive position just external to the margin of the dorsal

In this way are defined the limits of the future dorsal and ve lips, the former of which is distinguished by the presence of blood-vessel which runs along its entire margin. At the i (future posterior) angle of the nephrostome the vessel passes on to body of the nephridium and gradually extends backwards to f the nephridial vessel. This vessel also represents a space betw the peritoneum and the nephridial epithelium, and is formed a simple continuation of the nephrostomial vessel. A slight perito fold extending along the length of the nephridium (Plate 23, Fig. 26) marks in early stages the position of the future vessel. I more advanced nephridium represented in Fig. 27 the ves become well-defined in the anterior part of the nephridium, posteriorly it is still represented by the peritoneal fold. Late vessel completes its backward extension and forms, as well a the chief vascular supply of the glandular region and ter vesicle.

It is impossible in this place to give a full account of the m in which all the blood-vessels of *Arenicola* arise, and the brief account serves simply to explain the manner in whic characteristic relations arise between the segmental blood-vess the nephrostome. We have seen that the nephrostomial blood- is formed as a portion of the system of septal blood-vessels that its association with the funnel depends simply upon th that this structure itself represents in large part simply a differe part of the septum[1].

Shortly after the appearance of the system of septal bloo sels the membranous portion of the septum disappears[2]. nephrostome then undergoes a change of position, its dorsal assuming the antero-posterior direction of the nephridial bloo sel, and the aperture consequently becoming directed in

[1] For the vascular supply of the adult nephridium see GAMBLE & ASH 1900. Their Fig. 29 shows in a simple and clear manner the essential v relations of the nephridia. The detailed distribution of the blood-ve *A. cristata* has been the subject of investigation at the Marine Bi Laboratory, the results of which are not yet published.

[2] With the exception of a small portion extending from the anteri of the nephrostome to the body-wall and bearing the extra-nephridial the segmental vessel (see Plate 22, Fig. 6a, cf. also GAMBLE & ASHWORT Figs. 53—54).

te 25, Figs. 44, 46—49) as in the adult. The slip of transverse scle which binds the nephrostome down to the body-wall has already eared at the stage of Fig. 44, in a manner not satisfactorily erstood. The slip however is associated with the segmental d-vessel and is attached to the anterior border of the nephro- e near the point where the nephrostomial blood-vessel takes it

The fact of its attachment to the dorsal lip at the anterior e of the funnel becomes thus to some degree intelligible.

The origin of the germ-cells, which arise in connection with early nephridial vessels, may be briefly referred to here. Mere tion of the place and manner of origin of these cells will be cient, since the spermatogenesis and ovigenesis of *Arenicola* have the subject of special investigation at Woods Hole by other ors an account of whose work will it is hoped appear before long.

The nephridial vessels in somites VII—X (2nd to 5th nephridia) associated with the formation of the gonads. The early germ- ls in connection with each nephridium become distinguishable n after the appearance of the blood-vessel of the latter, and arise a proliferation of the peritoneal cells of its wall. They appear t on the anterior and first-formed portion of the vessel, i. e. in region immediately adjoining the posterior angle of the funnel, l at their first appearance (Plate 23, Fig. 27) have the form of newhat enlarged vesicular nuclei surrounded by a small quantity protoplasm and otherwise differing but slighty from the unaltered itoneal cells. As growth proceeds the germ-cells increase in nber and in size and acquire a very characteristic appearance te 25, Fig. 56). The nucleus becomes greatly enlarged and is. rrounded by a well-defined protoplasm of homogeneous appearance ;. 56 represents the appearance of these cells (oö-or spermatogonia) the second nephridium of a larva of a well-advanced stage. e large vesicular nuclei, whose chromatin is arranged in scattered mps closely applied to the nuclear membrane, already bear a se resemblance to the nuclei of the large oögonia or spermato- nia of the adult gonad. The germ-cells usually appear on their pective nephridia in the order of the formation of these organs, . in order from before back. I have never found them in con- tion with the first or last pair of nephridia, which typically do bear gonads in the adult.

The later development of the nephrostome consists essentially in extension of its epithelial surface, and in the completion of the

characteristic relations of this epithelium to the blood-vess
explained above, the formation of the series of dorsal vascul
cesses, and also of the vesicle of the ventral lip, is in all prob
primarily dependent upon the increase in epithelial surface. I
not however been able directly to study the early condition of
characteristic structures, which do not arise until a late st
development and have not yet begun their appearance in the
advanced larvae at my disposal.

Terminal Vesicle.

The terminal vesicle — as might have been expected fr
similarity of its adult structure to that of the glandular re
is formed as a differentiation of the most posterior portion
primitive nephridium. There is no ectodermal invaginati
from the resemblance in structure between the two regions it
appear a priori probable that the vesicle, like the tubular p
of the nephridium, is chiefly if not entirely of mesoblastic
A complete fusion, however, takes place at an early stage be
the ectoderm and the posterior end of the nephridium, and
from this region of fusion, in which the limits of ectoderm
mesoderm become indistinguishable, that the vesicle is differenti
In all probability, therefore, its distal portion is derived fr
region which originally was of ectoblastic origin; it seems how
impossible to decide the extent to which each germ-layer has t
part in its formation. In point of fact the vesicle is differenti
as a whole, without regard to limits of germ-layers, from the
treme posterior region where the nephridium and the ecto
become indistinguishably fused (see Plates 24 and 25, Figs.
37—42).

Fig. 34, Plate 24 represents the early nephridia of somites V a
VI in a larva of 14 somites. The nephridium is fused with the ec
derm slightly behind and internal to the ventral hooked setae a
point where the nephropore afterwards opens. At the point
fusion, ectoderm and mesoderm appear simply to become continu
with each other without any visible line of demarcation. The p
tion of the nephridium immediately adjoining the fusion rem
similar to the rest of the organ until a relatively late st
(cf. Plates 24 and 25, Figs. 37—42). Fig. 40 represents a cross sec
through the posterior portion of the first nephridium of a larva
the full number of somites and a length of about 1,75 mm.

of this region is vacolated and contains numerous coarse deeply-
ng granules, which are also present in the adjoining ectoderm;
o sharply defined terminal vesicle is as yet present. In the
idium represented in Fig. 42 (2nd nephridium of a somewhat
advanced larva) the conditions are essentially similar.
a Plate 25, Fig. 45, a representation of a cross section of a still
advanced stage, the earliest condition of the vesicle is seen.
erminal section of the nephridium has become dilated and has
ed thinner walls which however in all other respects are similar
se of the glandular regions proper. The strand of muscle-
which extends over the nephridium at the junction of the thick-
l glandular region represents a portion of the original longitu-
muscle-layer (which has already become associated with the
al region at the stage of Fig. 40). These muscle-fibres con-
the entire early musculature of the organ; later an extensive
rk of muscle-fibres is formed over the entire surface of the
le.
Figs. 52—55, Plate 25 represent a later condition of the vesicle,
has become much enlarged, and well defined with reference
adjoining glandular region. The most striking characteristic
vesicle in this larva was the presence of the large intensely
ng granules, both in the wall of the vesicle and in the neighboring
erm (see Fig. 51). The presence of granules in the ectoderm
s region is a common phenomenon, and is somewhat difficult
derstand. Possibly the granules in the preparation represent
of fluid excretory matters which find their way between the
ermal cells and collect there. No external opening at this
can as a rule be detected, and the contractions of the vesicle
possibly force the excretory matters into the intercellular spaces
ectoderm.
Fig. 6a, Plate 22 represents the vesicle at a later stage in a
ded condition, showing the thin walls composed of flattened
In larvae of this stage the contractions of the vesicle may
y be observed through the transparent body-walls. In later
opment the number of epithelial cells increases greatly and
ells themselves become columnar (see above p. 370). The
ulature of the wall also undergoes a great extension.

Literature.

This list includes papers and memoirs dealing with the structu development of the nephridia of Annelida. Reference has been omitted t papers that have been consulted relatively to the nephridia of other gro animals.

1. **Apáthy, St.,** Keimstreifen und Mesoblaststreifen bei Hirudineen. in: Z 14. Bd. 1891 pp. 388—393.
2. **Beddard, F. E.,** On the Occurrence of Numerous Nephridia in the Segment and on the Relationship between the Excretory Syste Annelida and in the Platyhelminths. in: Q. Journ. Micr. Sc. (2) V 1888 pp. 397—411 Pl. 30, 31.
3. —— On certain points in the Structure of *Urochaeta* and *Dichogast* further remarks on the Nephridia of Earthworms. Ibid. Vol. 2 pp. 235—282 Pl. 23, 24.
4. —— On the Homology between Genital Ducts and Nephridia in t chaeta. in: Proc. R. Soc. London Vol. 48 1890 pp. 452—455.
5. —— Researches into the Embryology of the Oligochaeta. 1. On points in the Development of *Acanthodrilus multiporus*. in: Q Micr. Soc. (2) Vol. 33 1892 pp. 497—540 Pl. 30, 31.
6. **Benham, W. Bl.,** The Nephridium of *Lumbricus* and its Blood with Remarks on the Nephridia in other Chaetopoda. Ibid. (2 1891 pp. 293—334 Pl. 22—25.
7. —— Postlarval Stage of *Arenicola marina*. in: Journ. Mar. Bi Plymouth Vol. 3 1893 pp. 48—52 Pl. 1.
8. **Bergh, R. S.,** Über die Metamorphose von *Nephelis*. in: Zeit. 41. Bd. 1884 pp. 284—301 Taf. 18, 19.
9. —— Die Exkretionsorgane der Würmer. in: Kosmos 17. Bd. 188 —122 Taf. 2.
10. —— Die Metamorphose von *Aulastoma gulo*. in: Arb. Z. Inst. W 7. Bd. 1885 pp. 231—291 Taf. 12—15.
11. —— Zur Bildungsgeschichte der Exkretionsorgane bei *Criodrilu* 8. Bd. 1888 pp. 223—248. Taf. 13, 14.
12. —— Neue Beiträge zur Embryologie der Anneliden. 1. Zur Ent und Differenzirung des Keimstreifens von *Lumbricus*. in: Zeit. 50. Bd. 1890 pp. 469—526 Taf. 19—21. 2. Die Schichtenbildung streifen der Hirudineen. Ibid. 52. Bd. 1891 pp. 1—17 Taf. 1, 2.
13. —— Nochmals über die Entwicklung der Segmentalorgane. Ibid 1899 pp. 435—449 Taf. 30.
14. **Bolsius, H.,** Recherches sur la structure des organes segmentaires dinées. in: Cellule Tome 5 1889 pp. 369—436 Pl. 1—3.
15. —— Nouvelles recherches sur la structure des organes segment Hirudinées. Ibid. Tome 7 1890 pp. 3—77 Pl. 1—3.
16. —— Les organes ciliées des Hirudinées (*Nephelis*). Ibid. 1891 pp. Pl. 1, 2.
17. **Bourne, A. G.,** On the structure of the Nephridia of the Medicina in: Q. Journ. Micr. Sc. (2) Vol. 20 1880 pp. 283—302 Pl. 24, 25.
18. —— Contributions to the Anatomy of the Hirudinea. Ibid. Vol. pp. 419—506 Pl. 24—34.

Bourne, A. G., Review: The Nephridia of Leeches. Ibid. Vol. 34 1894 pp. 545—565.

—— On certain points in the Anatomy and Development of some Earthworms. Ibid. Vol. 36 1894 pp. 11—33 Pl. 2—5.

Bürger, O., Beiträge zur Entwicklungsgeschichte der Hirudineen. Zur Embryologie von *Nephelis*. in: Z. Jahrb. Abth. Morph. 4. Bd. 1891 pp. 697—738 Taf. 41—43.

—— Neue Beiträge zur Entwicklungsgeschichte der Hirudineen. Zur Embryologie von *Hirudo medicinalis* und *Aulastoma gulo*. in: Zeit. Wiss. Z. 58. Bd. 1894 pp. 440—459 Taf. 26—28.

—— Weitere Beiträge zur Entwicklungsgeschichte der Hirudineen. Zur Embryologie von *Clepsine*. Ibid. 72. Bd. 1902 pp. 525—544 Taf. 30—32.

Child, C. M., The early Development of *Arenicola* and *Sternaspis*. in: Arch. Entwicklungsmech. 9. Bd. 1899 pp. 587—723 Pl. 21—25.

Cosmovici, L. C. E., Glandes génitales et organes segmentaires des Annélides polychètes. in: Arch. Z. Expér. Tome 8 1879—1880 pp. 233—372 Pl. 19—28.

Cunningham, J. T., On some points in the Anatomy of Polychaetes. in: Q. Journ. Micr. Sc. (2) Vol. 28 1887 pp. 229—278 Pl. 17—19.

Eisig, Hugo, Monographie der Capitelliden des Golfes von Neapel. in: Fauna Flora Golf. Neapel 16. Monogr. 1887.

Fauvel, Pierre, Les stades post-larvaires des Arénicoles. in: C. R. Acad. Sc. Paris Tome 127 1898 pp. 733—735.

Fraipont, J., Le genre *Polygordius*. in: Fauna Flora Golf. Neapel 14. Monogr. 1887.

Goodrich, Ed. S., On a new organ in the Lycoridea and on the Nephridium in *Nereis diversicolor*. in: Q. Journ. Micr. Sc. (2) Vol. 34 1893 pp. 387—402 Pl. 34, 35.

—— On the Coelom, Genital Ducts, and Nephridia. Ibid. Vol. 37 1895 pp. 477—510 Pl. 44, 45.

—— On the Nephridia of the Polychaeta. Part 1. On *Hesione*, *Tyrrhena* and *Nephthys*. Ibid. Vol. 40 1897 pp. 185—195 Pl. 6—9.

—— Id. Part 2. *Glycera* and *Goniada*. Ibid. Vol. 41 1898 pp. 439—457 Pl. 32—35.

—— On the Body-cavities and Nephridia of the Actinotrocha larva. Ibid. Vol. 47 1903 pp. 103—121 Pl. 8, 9.

Graf, Arnold, Beiträge zur Kenntnis der Exkretionsorgane von *Nephelis vulgaris*. in: Jena. Zeit. Naturw. (2) 21. Bd. 1893 pp. 163—195 Taf. 7—10.

—— Über den Ursprung des Pigments und der Zeichnung bei den Hirudineen. in: Z. Anz. 18. Bd. 1895 pp. 65—70.

—— Hirudineenstudien. in: Nova Acta Leop. Carol. 72. Bd. 1899 pp. 219—404 Taf. 1—15.

Hatschek, B., Studien über Entwicklungsgeschichte der Anneliden. in: Arb. Z. Inst. Wien 1. Bd. 1878 pp. 1—128 Taf. 1—8.

—— Über Entwicklungsgeschichte von *Echiurus*. Ibid. 3. Bd. 1881 pp. 1—34 Taf. 1—3.

—— Zur Entwicklung des Kopfes von *Polygordius*. Ibid. 6. Bd. 1886 pp. 1—12.

—— Entwicklung der Trochophora von *Eupomatus uncinatus* Phil. Ibid. 6. Bd. 1886 pp. 1—28 Taf. 1—5.

42. **Hoffmann, R. Wolfg.,** Beiträge zur Entwicklungsgeschichte der C chäten. in: Zeit. Wiss. Z. 66. Bd. 1899 pp. 335—357 Taf. 20, 21.
43. **Kleinenberg, N.,** Die Entstehung des Annelids aus der Larve von *Lo rhynchus*. Ibid. 44. Bd. 1886 pp. 1—227 Taf. 1—16.
44. **Kyle, H. M.,** Nephridia, Reproductive Organs, and Postlarval Stag *Arenicola*. in: Ann. Mag. N. H. (7) Vol. 18 1896 p. 295—301 Pl. 13—1
45. **Lang, Arnold,** Die Polycladen des Golfs von Neapel. in: Fauna Flora Neapel 11. Monogr. 1884.
46. —— Beiträge zu einer Trophocöl-Theorie. in: Jena. Zeit. Naturw. (2 38. 1903 pp. 1—376 Taf. 1—6.
47. **Maziarski, Stan.,** Recherches cytologiques sur les organes segmentaires Vers de terre. in: Poln. Arch. Biol. Med. Wiss. Lemberg 2. Bd. 1 pp. 1—81 Taf. 1—3.
48. **Mesnil, F.,** Les genres *Clymenides* et *Branchiomaldane* et les stade larvaires des Arénicoles. in: Z. Anz. 21. Bd. 1898 pp. 630—638.
49. **Meyer, Ed.,** Studien über den Körperbau der Anneliden. 1. in: M Stat. Neapel 7. Bd. 1887 pp. 592—741 Taf. 22—27. — 2. Ibid. 8. B pp. 462—662 Taf. 23—25. — 3. Ibid. 14. Bd. 1901 pp. 247—585 —17.
50. —— Die Abstammung der Anneliden. Der Ursprung der Metameri die Bedeutung des Mesoderms. in: Biol. Centralbl. 10. Bd. 1890 p —308.
51. **Moore, J. P.,** On the Structure of the Discodrilid Nephridium. in: J Morph. Boston Vol. 13 1897 pp. 327—380 Pl. 20—23.
52. **Rosa, D.,** Il cloragogo tipico degli Oligocheti. in: Mem. Accad. Sc. T (2) Tomo 52 1903 pp. 119—144.
53. —— Nefridii di Rotifero in giovani Lombrichi (Nota preliminare) Boll. Mus. Z. Anat. Comp. Torino Tomo 18 No. 440 1903.
54. **Schaeppi, Th.,** Das Chloragogen von *Ophelia radiata*. Eine morpholo physiologische Studie. in: Jena. Zeit. Naturw. (2) 28. Bd. 1894 p —293 Taf. 16—19.
55. **Vejdovský, Fr.,** Entwicklungsgeschichtliche Untersuchungen. Prag —1890.
56. —— Zur Entwicklungsgeschichte des Nephridialapparates von *Megas australis*. in: Arch. Mikr. Anat. 40. Bd. 1892 pp. 552—562 Taf. 32.
57. **Whitman, C. O.,** Embryology of *Clepsine*. in: Q. Journ. Micr. Sc. (2) V 1878 pp. 215—315 Pl. 12—15.
58. —— Germ-layers in *Clepsine*. in: Journ. Morph. Boston Vol. 1 1887 p —182 Pl. 4—6.
59. **Wilson, Edm. B.,** The Embryology of the Earthworm. Ibid. Vol. 3 pp. 387—462 Pl. 16—22.
60. **Woltereck, R.,** Trochophora-Studien. Über die Histologie der Larv die Entstehung des Annelids bei den *Polygordius*-Arten der No in: Zoologica Stuttgart 34. Heft 1902.

Explanation of Plates 22—25.

ıe figures of microscopic structure were drawn with the aid of the da under Zeiss objectives; Figs. 35 and 36 with 8 mm. obj., Oc. 6; ı 2 mm. apochromat., Oc. 1; Figs. 7, 8 and 29 with 2 mm. apochrom. the remainder with 2 mm. apochromat. Oc. 4. Finer details were wn in with Oc. 6.

Reference Letters.

e number of the somite is indicated by a Roman numeral. nephridium is designated by the letter n with an exponent to indi- ıber in the larval stage, thus: $n_2 = 2^{nd}$ larval nephridium (nephri- nite V) etc. The position of the septum is indicated by the numbers tes which it separates, thus: 5—6 = 5th septum.

vall.
ogen cells.
n-intestinal blood-sinus.
ılar muscle-fibres.
and second diaphragms.
ıl blood-vessel.
setae.
ıl seta-sacs.
elium of dorsal lip of nephro-
.
lerm of growing zone.
elium of ventral lip of nephro-
.
tory epithelium of nephri-
ory granules.
ells.
rmal gland-cells.
ındular region of nephri- ım.
g zone.
e.
mental septum.
e or excretophore.
ıl blood-vessel.
itudinal muscle-fibres.
ast cells.
ıchyme cells.
oderm pole-nucleus.
lar slip attached to dorsal nephrostome.

m.t.v Muscle-fibres of terminal vesicle.
m.v.l Membranous wall of vesicle of ventral lip of nephrostome.
n.b.v Nephridial blood-vessel.
nst.ep Epithelium of nephrostome.
nst.v Nephrostomial blood-vessel.
oes Oesophagus.
pb Proboscis.
pd Proctodaeum.
p.d.l Ciliated processes of dorsal lip of nephrostome.
p.m Post-mesodermal region of posterior extremity.
pn Peritoneum.
pn.f Peritoneal fold forming nephridial blood-vessel.
prost Prostomium.
s Septum.
sb Seta-forming ectodermal cell ('setiblast').
s.b.v Segmental blood-vessel.
s.oe.g Supra-oesophageal ganglion.
st Stomach.
t.m.f Transverse muscle-fibres.
t.v Terminal vesicle.
v.b.v Ventral or sub-intestinal blood-vessel.
v.b.w Ventral body-wall.
v.s Ventral seta.
v.s.s Ventral seta-sac.
v.v.l Vesicle of ventral lip of nephrostome.

Plate 22. Arenicola cristata.

Fig. 1. Portions of three successive adult nephridia, showing gene ance after exposure by removal of transverse muscles. X

Fig. 2. First adult nephridium of left side, with nephrostome turn and spread open so as to show structure of dorsal and v × 7.

Fig. 3. Second adult nephridium of left side, seen in its natural po above; showing nephrostome, with slip of transverse musc to the anterior border of its dorsal lip; also position of ne blood-vessel and gonad. × 5.

Fig. 4. Fourth nephridium of left side with nephrostome freed fro ment and turned outward so as to show the outer po ventral lip. × 5.

Fig. 5. Portion of transverse section of wall of glandular portio dium. Explanation in the text. × 800.

Fig. 6. Portion of transverse section of epithelial wall of the vesi lip of nephrostome, showing the abrupt transition to the portion of the same. × 800.

Fig. 6a. Longitudinal section of first nephridium of a larva of 5- tangentially to body-wall, showing terminal vesicle, me necting nephrostome to body-wall, and nephridial blood- structure of the excretory cells. Section passes below stome. × 800.

Fig. 7. Sagittal section through a larva of the swarming stag in process of formation), showing general characteristics this stage. Nerve cord differentiated in ventral ecto incomplete dorsally; early lumen of stomach indicated; the extreme posterior region in process of differentiation

Fig. 8. Sagittal section through larva at beginning of crawling s VI in process of formation), showing early septa, growi early proctodaeum. Absorption of yolk has progressed since stage of Fig. 7. × 800.

Fig. 9. Horizontal section through posterior region of larva in whi is in process of formation, showing early differentiation o structures, septa, ventral blood-vessel, chloragogen cell × 800.

Plate 23. Arenicola cristata.

Fig. 10. Oblique section through posterior region of larva in whi is in process of formation, showing large posterior meso growing zone, early septa, early nephridial rudiments blood-vessel. × 800.

Fig. 11. Longitudinal section through body-wall of a larva in whic is in process of formation, showing early pronephridia o and 5. Goblet-cells in ectoderm also shown. × 800.

Fig. 12. Cross section through ventral portion of intersegmental of a larva in which somite VII is just appearing, showing la cells all similar to one another in appearance. × 800.

Cross section through intersegmental region 5—6 of a larva in which somite VII is in process of formation, showing disposition and histological structure of mesoblast in region of differentiating 6th septum. Longitudinal muscle-fibres appearing. × 800.

Transverse section through growing zone of a larva in which somite VI is in process of formation. Section passes somewhat obliquely forwards from above down. Dorsally the undifferentiated mesoblast is shown; ventrally the region of formation of early 5th septum and longitudinal muscle-fibres. Ectodermal cells large and well defined with large nuclei. × 800.

Cross section through intersegmental region 5—6 in a larva in which somite VII is just appearing; showing on the right side the commencing differentiation of the third larval nephridium (somite VI); lumen just appearing in cells otherwise similar to the adjoining myoblast cells. Ectoderm cells well defined. On left side is shown early condition of dorsal and ventral seta-sacs. × 800.

Cross section through anterior of somite IV of a larva in which somite IX is just appearing. 1st provisional nephridium shown in cross section. Myoblast cells reduced. × 800.

Similar cross section through somite V of the same larva. × 800.

Similar cross section through somite VI of same larva. × 800.

Similar cross section through somite VII of same larva. On left side nephridium further advanced than on right. Muscle-fibres less developed than in preceding somites, and myoblast cells larger. × 800.

Similar cross-section through somite VIII of the same larva. On the left the section passes more anteriorly than on right and there shows intersegmental region 7—8 with myoblasts. Nephridal cell not yet distinguishable from the others. Shows also early seta-sacs which still retain connection with the ectoderm and to which flattened mesoderm cells are applied. Muscle-fibres less developed than in preceding somite. × 800.

Section through region of formation of somite IX of the same larva Ventrally the section passes through region of formation of 8th septum; dorsally through the anterior border of the pigmented post-mesodermal region. Large well-defined cells of the ectodermal growing zone. Longitudinal muscle-fibres just making their appearance. × 800.

Cross section through post-mesodermal region of same larva, showing small, pigmented, vacuolated, radially arranged ectoderm cells with their small deeply staining nuclei. A few flattened mesoderm cells applied to body-wall and proctodaeum. Strong cilia on ventral wall of proctodaeum. × 800.

Cross section through anterior region of somite XI of a larva in which somite XII is in process of formation; showing commencing differentiation of 8th larval nephridium (6th definitive). Nephridium somewhat more advanced in development on the left side. Myoblast cells well defined but apparently undergoing reduction. Large embryonic nuclei in ectoderm; smaller nuclei in the ventral region have the characters of tissue nuclei, and are probably connected with the differentiating nerve-cord. × 800.

Fig. 24. Cross section through intersegmental region 11—12 of a larva i somite XII is appearing; showing early mesoblast; intestine surr by blood-sinus continous with dorsal and ventral blood-vessels well-defined ectoderm cells of the growing zone. Differentia the ventral nerve-cord is in progress, and smaller, more deeply s nuclei are seen in its vicinity. × 800.

Fig. 25. Cross section through nephridium and adjacent body-wall of so of a larva of ca. 1.5 mm. length and ca. 45 somites. Nephridi definitive of right side) still apparently one cell in cross s showing also a peritoneal fold running along nephridium from in a position corresponding to the later nephridial blood Nucleus large and nucleolated. Protoplasm vacuolated and cor granules; stains more deeply at border of lumen. Cross-c visible in lumen. × 800.

Fig. 26. Cross section through nephridium and adjacent body-wall of som of a larva of a later stage than that of Fig. 25 (ca. 1.6 mm. somites). Nephridium (3rd definitive of left side) has become into cells and lumen has become intercellular; 3 cells present section at this region. Histological characteristics of cells sam Fig. 25. A similar membrane occupies position of future ne blood-vessel.

Fig. 27. Cross section through nephridium and adjacent body-wall of som (3rd definitive nephridium, left side) of a more advanced lar that of Fig. 26 (ca. 2,0 mm.; full number somites); section pas behind nephrostome and shows nephridial blood-vessel with a germ cell applied to it. × 800.

Fig. 28. Similar cross section through middle region of 1st definitive dium (somite VI) of right side of a larva of later stage than Fig. 27 (2.5 mm.; full number somites). Nephridium 5 cells i section; histological structure as above. Nephridial blood present in anterior portion of nephridium. × 800.

Fig. 29. Horizontal section through posterior body-region of a larva in somite XI is being formed. Section passes along level of ne and ventral setae, and shows the entire series of early ne × 400.

Fig. 30. Sagittal section through posterior body-region of a larva in somite XI is being formed. Shows early septa, intestine, nerv Septa in advance of somite VIII are dorsally incomplete. Sho large terminal mesoblast-nucleus. × 800.

Fig. 31. Longitudinal section through early region of formation of ne in a larva in which somite XI is just appearing; showing early and associated nephridial rudiments of the 6th and 7th larval r dia (somites IX and X). Pole-nucleus of mesoderm and mitosis differentiated mesoderm cells also seen. Also the large ecto nuclei of the growing zone. × 800.

Plate 24. Arenicola cristata.

Fig. 32. Nearly horizontal section through the posterior region of a la which somite XVI is in process of formation. The ectodermal g

zone, and the differentiation of the seta-sacs and early septa are shown. The proctodaeal cilia in the post-mesodermal region are shown; also the mesodermal nuclei applied to the proctodaeum and body-wall of this region. × 800.

33. Horizontal section through the posterior achaetous region of a more advanced larva (ca. 1.4 mm. length; ca. 45 somites) showing the growing zone, and formation of early septa of this region. Post-mesodermal region sharply defined; a valve separating proctodaeum and rectum; circular muscle-fibres applied to body-wall and anal region in this region. × 800.

34. Part of a horizontal section of a larva in which somite XIV is appearing. Section passes tangentially to median ventral region a little to left of nerve-cord. Shows early nephridia of somites V and VI (2nd provisional and 1st definitive nephridia), and their relations to septa, ventral hooked setae and body-wall. × 800.

35. Sagittal section of a larva in which somite VIII is forming; showing early septa, incomplete dorsally. First septum distinctly separate from retractor muscle of proboscis. Diaphragms have not yet appeared. × 300.

36. Sagittal section of a larva in which somite XIV is forming; showing the series of septa and early stage of 2nd diaphragm. Remains of 3rd primitive septum seen stretching from insertion of diaphragm to anterior region of stomach. × 300.

37. Horizontal section of left nephridium of somite VI (1st definitive) in larva of ca. 2 mm. and 50 somites, showing early condition of nephrostome, and its continuity with the septum. × 800.

38. Portion of a horizontal section through a more advanced larva than the preceding (1.5 mm. long; 50 + somites); showing nephridium of somite IX (fourth definitive), nerve-cord, and body-wall, cut tangentially. Posteriorly the nephridium is fused with the ectoderm; no terminal vesicle is as yet differentiated. Continuity of the nephrostomial lip with the primitive septum, and connection of the latter with the ventral blood-vessel, are visible. The smaller cells at the anterior end of the nephridium form the early nephrostomial epithelium. × 800.

39. Sagittal section through nephridium and adjoining body-wall of somite VI (1st definitive nephridium) of a larva of about same stage as last (ca. 45 somites; ca. 1.7 mm. long). Showing continuity of septum with dorsal lip of early nephrostome. The most anterior cells of the nephridium are small and cubical; closely crowded nuclei in the portion of the septum immediately continuous with the dorsal lip probably indicate a stage in the transformation of this portion of the septum into funnel-cells. No differentiated terminal vesicle as yet. Nephridial blood-vessel not yet formed. × 800.

40. Cross section through posterior extremity of 1st nephridium (somite VI) of a larva of a more advanced stage than that of Fig. 38 (ca. 2.25 mm.; full number of somites), showing the posterior continuity of nephridium and ectoderm and the presence of excretory granules in the latter. No differentiated terminal vesicle as yet. The terminal region is overarched by a band of muscle-fibres continuous with the longitudinal

fibres of the body-wall. These represent the first appearance of the musculature of the terminal vesicle (compare Figs. 45, 53, 55). × 800.

Fig. 41. Horizontal section through ventral region of somites VII and VIII of a more advanced larva (ca. 3 mm.; full number of somites), showing the origin of the segmental vessels from the ventral blood-vessel, and the relations of one of the vessels to the 2nd nephridium of the right side. × 200.

Plate 25. Arenicola cristata.

Fig. 42. Longitudinal section through the left nephridium of somite VII (2nd definitive nephridium) and adjacent body-wall of the larva of Fig. 41, showing (1) a portion of the nephridial blood-vessel to which is applied a large germ-nucleus; (2) the posterior junction of the nephridium with the ectoderm (corresponding to the early undifferentiated terminal vesicle); and (3) the cubical cells of the early nephrostome with their shorter cilia. The lumen has extended throughout the entire length of the organ and seems to communicate with the exterior; the external opening, however, is very minute. × 800.

Fig. 43. Cross section through the anterior end of the first left definitive nephridium (somite VI) of a larva of about the same stage as Fig. 42, showing the early condition of the septal blood-vessels of this somite, and their relations to the nephridia and to the body-wall. Portions of the primitive septum still remain. × 800.

Fig. 44. Longitudinal section through the anterior part of the 4th nephridium (somite IX) of a larva of a somewhat more advanced stage than Fig. 42 (length ca. 3 mm.). The nephrostome is turned somewhat inward and has been cut in such a way as to show its epithelium in cross section; the figure shows also its relations to the nephrostomial blood-vessel, which runs across its dorsal lip and is in close relation to the transverse band of muscle (*t. m. f.*), which is also attached to this lip and binds it down to the ventral body-wall. A portion of the primitive septum in which the blood-vessel appears is seen at *s*. × 800.

Fig. 45. Cross section through the posterior end of the 1st nephridium (somite VI) of a larva in which the terminal vesicle is beginning to differentiate (length ca. 3.5 mm; full no. of somites). The terminal portion adjoining the ectoderm is thin walled and somewhat dilated. A band of muscle-fibre (derived from the longitudinal layer of the body-wall) extends across the terminal vesicle at its junction with the glandular region proper. Granules in the ectoderm near the terminal vesicle. × 800.

Figs. 46-50. These figures represent successive transverse sections, 7.5 μ thick, through the nephrostome and anterior glandular portion of the first nephridium (somite VI) of a larva of ca. 4 mm. in length. The nephrostome has become well-defined and is composed of a single layer of small cubical ciliated cells; along its dorsal lip runs the nephrostomial blood-vessel. The continuity of this vessel with the nephridial vessel is shown. The relation of the segmental blood-vessel to the transverse muscle-band attached to the dorsal lip is also shown. *l*, vacuolated wandering cells (excretophores?). × 800.

Figs. 51-54. These figures similarly represent four successive sections of the terminal vesicle of the same nephridium. Large deeply staining granules are present in the walls of the vesicle, and in the adjacent ectoderm of the body-wall. × 800.

Fig. 55. Cross section of the terminal vesicle and adjacent glandular region of the 2nd nephridium of the left side (somite VII) of the same larva. The flattened cells of the vesicle with their contained excretory granules are shown. Also the early muscle-band crossing the vesicle at its junction with the glandular region. × 800.

Fig. 56. Cross section through the anterior region of the nephridium of Fig. 55, immediately behind the funnel, showing the nephridial blood-vessel, with well-defined germ-cells (rudimentary ovary or testis) applied to it. × 800.


5.
8.
prost
mth.
pb.
s.
g.z.
II
III
IV
V
VI
9.
pd.
gl.c.
gl.c.
s.oeg.
VII
g.z.
VI
n3
i.s.
V
I
n2
IV
n1
v.b.v.
c.c.
st.



Lith. Anst. v. E. A. Funke, Leipzig.

39.


c.m.f.
ect.
v.s.
v.s.s.
l.m.f.
n.st.ep.
st.
n.st.ep.
l.
s.
v.b.v.
c.c.

34.


IV
l.m.f.
ex.g.
v.b.w.
l.m.f.

41.


v.s.
s.b.v. (sV-VI)
n.b.v.
v.b.v.
c.c.